



Xiphosurid fossils from the Pennsylvanian Beeman Formation, Otero County, New Mexico

Spencer G. Lucas, Allan J. Lerner, William A. Dimchele, Amanda K. Cantrell, Thomas L. Suazo, and Dan S. Chaney, 2014, pp. 311-314

in:

Geology of the Sacramento Mountains Region, Rawling, Geoffrey; McLemore, Virginia T.; Timmons, Stacy; Dunbar, Nelia; [eds.], New Mexico Geological Society 65th Annual Fall Field Conference Guidebook, 318 p.

This is one of many related papers that were included in the 2014 NMGS Fall Field Conference Guidebook.

Annual NMGS Fall Field Conference Guidebooks

Every fall since 1950, the New Mexico Geological Society (NMGS) has held an annual [Fall Field Conference](#) that explores some region of New Mexico (or surrounding states). Always well attended, these conferences provide a guidebook to participants. Besides detailed road logs, the guidebooks contain many well written, edited, and peer-reviewed geoscience papers. These books have set the national standard for geologic guidebooks and are an essential geologic reference for anyone working in or around New Mexico.

Free Downloads

NMGS has decided to make peer-reviewed papers from our Fall Field Conference guidebooks available for free download. Non-members will have access to guidebook papers two years after publication. Members have access to all papers. This is in keeping with our mission of promoting interest, research, and cooperation regarding geology in New Mexico. However, guidebook sales represent a significant proportion of our operating budget. Therefore, only *research papers* are available for download. *Road logs, mini-papers, maps, stratigraphic charts*, and other selected content are available only in the printed guidebooks.

Copyright Information

Publications of the New Mexico Geological Society, printed and electronic, are protected by the copyright laws of the United States. No material from the NMGS website, or printed and electronic publications, may be reprinted or redistributed without NMGS permission. Contact us for permission to reprint portions of any of our publications.

One printed copy of any materials from the NMGS website or our print and electronic publications may be made for individual use without our permission. Teachers and students may make unlimited copies for educational use. Any other use of these materials requires explicit permission.

This page is intentionally left blank to maintain order of facing pages.

XIPHOSURID FOSSILS FROM THE PENNSYLVANIAN BEEMAN FORMATION, OTERO COUNTY, NEW MEXICO

SPENCER G. LUCAS¹, ALLAN J. LERNER¹, WILLIAM A. DIMICHELE², AMANDA K. CANTRELL¹,
THOMAS L. SUAZO¹, AND DAN S. CHANEY²

¹ New Mexico Museum of Natural History, Albuquerque, NM, spencer.lucas@state.nm.us

² Department of Paleobiology, NMNH Smithsonian Institution, Washington, D.C.

ABSTRACT—We document body fossils of the xiphosurid (horseshoe crab) *Euproops danae* Meek and Worthen from lacustrine black shale of the Missourian interval of the Beeman Formation near Alamogordo, New Mexico. These xiphosauroids, associated with conchostracans, bivalves, microconchids, fish scales and a paleoflora, are from a likely freshwater lake deposit. This is the first report of *E. danae* from the western USA and the first documentation of xiphosurid body fossils in the Pennsylvanian of New Mexico.

INTRODUCTION

Xiphosurids (Order Xiphosura), the true horseshoe crabs, are chelicerate arthropods with a fossil record that goes back to the Late Ordovician (Ashgillian, ~445 Ma) of central Canada (Rudkin et al., 2008). They are generally considered to be a sister group to eurypterids and arachnids within the Euchelicerata clade, although other alternatives have also been proposed (Dunlop, 2010). Their body and trace fossils are particularly well known from Carboniferous and Permian strata in North America and Europe (e.g., Filipiak and Krawczyński, 1996; Babcock and Merriam, 2000).

In New Mexico, the only late Paleozoic record of xiphosurids is of their trace fossils, notably their well-known walking traces (trackways) assigned to the ichnogenus *Kouphichnium* (e.g., Minter and Braddy, 2009). We document xiphosurid body fossils discovered in Pennsylvanian strata in the Sacramento Mountains of Otero County, New Mexico (Fig. 1). These specimens are the first occurrence of Pennsylvanian xiphosurid body fossils known from New Mexico.

PROVENANCE

The xiphosurid fossils documented here were collected at NMMNH (New Mexico Museum of Natural History, Albuquerque) locality 9109 in the Pennsylvanian Beeman Formation in La Luz Canyon, northeast of Alamogordo (Fig. 1). These fossils came from an approximately 2-m-thick interval of black shale at the base of a 10 m-thick, coarsening upward lake-fill deposit. The deposit also yields conchostracans, bivalves, microconchids, fish scales and a paleoflora. The floral elements are the most abundant fossils in the deposit. The flora is dominated by plants typical of seasonally dry climatic conditions, including common cordaitalean foliage with a wide variety of less common elements including *Sphenopteridium*, *Charliea*, *Taeniopteris* and walchian conifers; the flora also includes uncommon taxa typical of wet substrates—*Pecopteris* tree ferns, the pteridosperms *Alethopteris* sp., *Neuropteris* sp., and cf. *Macroneuropteris scheuchzeri*, and the calamitalean foliage *Annularia* cf. *spinulosa* and a small form similar to *A. asteris*, which probably lived

only along the margins of the water body. The stratigraphic section that includes this deposit is in the middle part of the Beeman Formation and thus is of Missourian age (Pray, 1961; Raatz and Simo, 1998).

DESCRIPTION AND IDENTIFICATION

Two xiphosurid body fossils were collected at NMMNH locality 9109 and are described here. NMMNH P-67802 is a part/counterpart of an opisthosoma (thoracetron) in dorsal aspect (Fig. 2A-B). The opisthosoma is broad and rounded in appearance, with a width of ~16 mm. It consists of an axial column, pleural lobes and a marginal rim. The axial column is ~4 mm wide with a preserved length of 8 mm. It has a prominent posterior lobe containing a large tubercle at its center. The tubercle has at its anterior border a short, anteriorly directed spine. The lobe has sharply angled lateral borders that give it a sub-triangular appearance. The telson is not preserved. The axial column is missing the anterior first and second rings. There is a median axial node on the third ring. The third through fifth rings also contain paired lateral bosses. The pleural lobes are divided into segments by transverse ribs, some of which extend onto the marginal rim and appear as spines. The opisthosomal rim is ~2 mm wide and shows a scalloped border particularly at its posterior.

The second specimen, NMMNH P-67801, is also a part/counterpart of an opisthosoma (thoracetron) in dorsal aspect (Fig. 2C). It is comparable in morphology to P-67802, but not as well preserved. It too has a width of ~16 mm. There is a small (~2 mm) molluscan shell preserved midway on the right border of the axial column.

These specimens, although incomplete, show diagnostic features that justify assignment to *Euproops danae* Meek and Worthen, 1865, including a median tubercle on the third ring of the axial column, a spined tubercle on the posterior lobe of the axial column and pleural ridges that end in marginal spines (e.g., Størmer, 1955; Ambrose and Romano, 1972; Anderson, 1994; Anderson and Selden, 1997; Babcock and Merriam, 2000). We note, however, that the genus *Euproops* is currently under review, with the possibility being that supposed separate species

may represent various ontogenetic stages within a single species (Haug et al., 2012).

DISCUSSION

Euproops danae is a xiphosurid previously documented from Illinois (Fisher, 1977, 1979), Indiana (Raymond, 1944), Kansas (Babcock and Merriam, 2000), Ohio (Murphy, 1970), Pennsylvania (Raymond, 1944), Nova Scotia (Copeland, 1957), Great Britain (Ambrose and Romano, 1972; Anderson 1994), Germany (Brauckmann, 1982) and some other European locations (France, Czech Republic, Ukraine: J. Schneider, written commun., 2014). Its New Mexico record documented here thus appears to be the first occurrence of *E. danae* in the western USA.

The paleoecology of the site that yielded the xiphosurid fossils, in a broad sense, is best indicated by the combination of the sedimentary environment and the fossil flora. The shale and siltstone layers from which the animal and plant fossils are derived comprise a 10-meter thick coarsening upward

sequence. At the base, a dark, organic-rich shale rests directly on medium-grained, ripple laminated sandstone. The upper parts of the basal sandstone contain foliage of the coniferophyte *Cordaite*s, and there are thin layers enriched in fine-grained sediment that are also mudcracked. The dark shale is clay rich and contains a well preserved flora dominated by cordaitalean foliage, with a moderately diverse assemblage of subsidiary taxa, possibly washed into the deposits, the most common of which mirror *Cordaite*s in being typical of seasonally dry environments: *Walchia*, *Sphenopteridium*, *Charliea* and *Taeniopteris*. This flora includes a small mixture of more typically wetland plants, such as the pteridosperms *Neuropteris*, *Macroneuropteris*, and *Alethopteris*, calamitalean foliage, and the tree fern foliage, *Pecopteris*. The flora changes little upward, becoming increasingly fragmentary and allochthonous in character as the sediment becomes both coarser and less organic rich. The deposit terminates in a thin, 3-cm-thick coaly layer that is overlain by 25 cm of ripple-bedded sandstone.

The thinly laminated shale (green to black in color) also contains a low-diversity fauna composed of bivalves, microconchids (spirorbids), and conchostracans, in addition to the xiphosurids. This fauna is not diagnostic of a particular salinity level, although it suggests that conditions were certainly not marine, though the presence of xiphosurids suggests a connection to a marine environment. These animals would be consistent with slight salinity to freshwater. Conchostracans, in particular, are consistent with conditions of intermittent dryness in the surrounding landscape.

There is insufficient lateral exposure to characterize the geometry of this deposit, but the general features – a basal shallow-water, ripple laminated sandstone with evidence of subaerial exposure, suggests a channel with intermittent periods of low water. This channel was abandoned and flooded and may have become a lake, indicated by the organic shales at the base. Its proximity to marine strata (above and below) and the presence of xiphosurids suggests this was a coastal lake. Subsequent filling ensued and a swampy habitat developed at this site near as the former channel filled.

The mixture of plants typical of seasonally dry environments with those typical of wet substrates is characteristic of many Pennsylvanian floras described from the western parts of the US, then equatorial Pangea (see review in DiMichele, 2014). Dark, organic-rich shales containing floras and faunas similar to the one described here have been reported elsewhere from the Missourian of central New Mexico (e.g., Lerner et al., 2009), where they also appear to represent small lakes, set in a background of climatic seasonality. This strongly differentiates the western regions of Pangea from the more central portions of the supercontinent (Tabor and Poulsen, 2008), where the wetter portions of glacial-interglacial cycles were marked by the formation of coal beds.

Euproops danae is a characteristic Carboniferous coal swamp form that is usually associated with freshwater habitats (e.g., Babcock and Merriam, 2000). The first published descriptions of *E. danae* were from the well known Pennsylvanian (Desmoinesian) Mazon Creek deposits of Illinois (Meek and




FIGURE 1. Late Paleozoic stratigraphy in the Sacramento Mountains (after Raatz and Simo, 1998) showing stratigraphic position of xiphosurid specimens (drawing of *Euproops*, after Anderson, 1994) in Beeman Formation.




FIGURE 2. Specimens of *Euproops danae* from the Beeman Formation near Alamogordo, New Mexico. A-B, NMMNH P-67802, part and counterpart. C, NMMNH P-67801, part. Scale bars = 5 mm.

Worthen, 1865; Meek, 1867a, b) where they are preserved in siderite concretions. Its occurrence in Mazon Creek deposits is restricted to the freshwater Braidwood fauna, although it is sometimes found as fragmentary inclusions within coprolites of the marine Essex fauna (Mikulic, 1998). Preservation bias for *Euproops danae* being restricted to the Braidwood fauna can be excluded due to preservation being similar in both the Braidwood and Essex assemblages (Gray, 1988). Fisher (1979) suggested that *Euproops danae* was capable of subaerial excursions, although this was largely countered by Anderson (1994).

Xiphosurids in general have an overall poor body fossil record, which is due in part to their having a non-mineralized exoskeleton. There is also a poor representation in the fossil record of the depositional environments in which most fossil xiphosurids occur (Babcock et al., 2000). The New Mexico *Euproops* specimens appear somewhat unusual in regard to their mode of preservation as bedding plane compressions. *Euproops* is more typically seen preserved within concretions. The New Mexico specimens share comparable thoracetrone widths, indicating that both individuals were of the same ontogenetic stage. Their fragmentary condition, however, makes it difficult to determine whether they were juveniles or adults, though they are likely adults based on the size data in Copeland (1957). It is possible that the New Mexico specimens are incomplete molts rather than having originated from dead individuals. Continued collecting within the Beeman Formation will hopefully provide more complete specimens.

ACKNOWLEDGMENTS

Joerg Schneider and Matt Stimson provided helpful reviews of the manuscript.

REFERENCES

- Ambrose, T. and Romano, M., 1972, New Upper Carboniferous Chelicerata (Arthropoda) from Somerset, England: *Palaeontology*, v.15, p. 569–578.
- Anderson, L.I., 1994, Xiphosurans from the Westphalian D of the Radstock Basin, Somerset Coalfield, the South Wales Coalfield and Mazon Creek, Illinois: *Proceedings of the Geologists Association*, v.105, p. 265–275.
- Anderson, L.I. and Selden, P.A., 1997, Opisthosomal fusion and phylogeny of Palaeozoic Xiphosura: *Lethaia*, v.30, p. 19–31.
- Babcock, L.E. and Merriam, D.F., 2000, Horseshoe crabs (Arthropoda: Xiphosuridae) from the Pennsylvanian of Kansas and elsewhere: *Transactions of the Kansas Academy of Science*, v. 103, p. 76–94.
- Babcock, L.E., Merriam, D.F. and West, R., 2000, *Paleolimulus*, an early limuline (Xiphosurida), from Pennsylvanian-Permian Lagerstätten of Kansas, and taphonomic comparison with modern *Limulus*: *Lethaia*, v. 33, p. 129–141.
- Brauckmann, C., 1982, Der Schwertschwanz *Euproops* (Xiphosurida Limulina, Euproopacea) aus dem Ober-Karbon des Piesbergs bei Osnabrück: *Osnabrück. Naturwissenschaftlichen Mitteilungen*, v. 9, p.17–26.
- Copeland, M.J., 1957, The arthropod fauna of the Upper Carboniferous rocks of the Maritime Provinces: *Geological Survey Canada Memoir*, v. 286, 110 p.
- DiMichele, W.A., 2014, Wetland-dryland vegetational dynamics in the Pennsylvanian ice-age tropics: *International Journal of Plant Sciences*, v. 175, p. 123–164.

- Dunlop, J.A., 2010, Geological history and phylogeny of Chelicerata: Arthropod Structure and Development, v. 39, p. 124–142.
- Filipiak, P., and Krawczyński, W., 1996, Westphalian xiphosurans (Chelicerata) from the Upper Silesia Coal Basins of Sosnowiec, Poland: Acta Palaeontologica Polonica, v. 41, 4, p. 413–425.
- Fisher, D.C., 1977, Functional morphology of spines in the Pennsylvanian horseshoe crab *Euproops danae*: Paleobiology, v.3, p. 175–195.
- Fisher, D.C., 1979, Evidence for subaerial activity of *Euproops danae* (Merostomata, Xiphosurida); in Nitecki, M.H., ed., Mazon Creek fossils: Academic Press, New York, p. 379–447.
- Gray, J., 1988, Evolution of the freshwater ecosystem: The fossil record: Palaeogeography, Palaeoclimatology, Paleoecology, v. 62, p. 1–214.
- Haug, C. Van Roy, P., Leipner, A., Funch, P., Rudkin, D.M., Schöllmann and Haug, J. T., 2012, A holomorph approach to xiphosuran evolution—a case study on the ontogeny of *Euproops*: Developmental Genes and Evolution, v. 222, p. 253–268.
- Lerner, A.J., Lucas, S.G., Spielmann, J.A., Krainer, K., DiMichele, W.A., Chaney, D.S., Schneider, J.W., Nelson, W.J. and Ivanov, A., 2009, The biota and paleoecology of the Upper Pennsylvanian (Missourian) Tinajas locality, Socorro County, New Mexico: New Mexico Geological Society, Guidebook 60, p. 267–280.
- Meek, F.B., 1867a, Note on *Bellinurus danae*, from the Illinois Coal Measures: American Journal of Science, 2nd Series, v. 43, p. 257–258.
- Meek, F.B., 1867b, Note on a new genus of fossil Crustacea: American Journal of Science, 2nd Series, v. 43, p. 394–395.
- Meek, F.B., and Worthen, A.H., 1865, Notice of some new types of organic remains, from the Coal Measures of Illinois: Proceedings of the Academy of Natural Sciences, v. 17, p. 41–53.
- Mikulic, D.G., 1998, Xiphosura, in Shabica, C.W. and Hays, A.A., eds., Richardson's guide to the fossil fauna of Mazon Creek: Northeastern Illinois University Press, Chicago, p. 134–139.
- Minter, N.J. and Braddy, S.J., 2009, Ichthyology of an Early Permian intertidal flat: The Robledo Mountains Formation of southern New Mexico, USA: Special Papers in Paleontology, no.82, 107 p.
- Murphy, J.L., 1970, *Euproops* from the “Uffington Shale” of Columbiana County, Ohio: Annals of the Carnegie Museum, v. 41, p. 281–286.
- Pray, L.C., 1961, Geology of the Sacramento Mountains escarpment, Otero County, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Bulletin 35, 144 p.
- Raatz, W.D. and Simo, J.A., 1998, The Beeman Formation (Upper Pennsylvanian) of the Sacramento Mountains, New Mexico: Guide to the Dry Canyon area with discussion of shelf and basin responses to eustasy, tectonics, and climate: New Mexico Geologic Society, Guidebook 49, p. 161–176.
- Raymond, P.E., 1944, Late Paleozoic xiphosurans: Museum of Comparative Zoology Bulletin, no. 94, p. 475–508.
- Rudkin, D.M., Young, G.A. and Nowlan, G.S., 2008, The oldest horseshoe crab: A new xiphosurid from Late Ordovician Konservat-Lagerstätten deposits, Manitoba, Canada: Palaeontology, v. 51, p. 1–9.
- Störmer, L., 1955, Merostomata; in Moore, R.C., ed., Treatise on invertebrate paleontology, P, Arthropoda 2: Geological Society of America and University of Kansas, Boulder and Lawrence, p. 4–175.
- Tabor, N.J. and Poulsen, C.J., 2008, Palaeoclimate across the Late Pennsylvanian–Early Permian tropical palaeolatitudes: a review of climate indicators, their distribution, and relation to palaeophysiographic climate factors: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 268, p. 293–310.



Anthropogenic “clast” in stream sediment in Sawmill Canyon east of Mayhill. Photo courtesy of Ben Hallett.