

New Mexico Geological Society

Downloaded from: <https://nmgs.nmt.edu/publications/guidebooks/21>



Oil and gas possibilities in the Pedregosa basin

Eugene Greenwood

1970, pp. 105. <https://doi.org/10.56577/FFC-21.105>

in:

Tyrone, Big Hatchet Mountain, Florida Mountains Region, Woodward, L. A.; [ed.], New Mexico Geological Society 21st Annual Fall Field Conference Guidebook, 176 p. <https://doi.org/10.56577/FFC-21>

This is one of many related papers that were included in the 1970 NMGS Fall Field Conference Guidebook.

Annual NMGS Fall Field Conference Guidebooks

Every fall since 1950, the New Mexico Geological Society (NMGS) has held an annual [Fall Field Conference](#) that explores some region of New Mexico (or surrounding states). Always well attended, these conferences provide a guidebook to participants. Besides detailed road logs, the guidebooks contain many well written, edited, and peer-reviewed geoscience papers. These books have set the national standard for geologic guidebooks and are an essential geologic reference for anyone working in or around New Mexico.

Free Downloads

NMGS has decided to make peer-reviewed papers from our Fall Field Conference guidebooks available for free download. This is in keeping with our mission of promoting interest, research, and cooperation regarding geology in New Mexico. However, guidebook sales represent a significant proportion of our operating budget. Therefore, only *research papers* are available for download. *Road logs*, *mini-papers*, and other selected content are available only in print for recent guidebooks.

Copyright Information

Publications of the New Mexico Geological Society, printed and electronic, are protected by the copyright laws of the United States. No material from the NMGS website, or printed and electronic publications, may be reprinted or redistributed without NMGS permission. Contact us for permission to reprint portions of any of our publications.

One printed copy of any materials from the NMGS website or our print and electronic publications may be made for individual use without our permission. Teachers and students may make unlimited copies for educational use. Any other use of these materials requires explicit permission.

This page is intentionally left blank to maintain order of facing pages.

OIL AND GAS POSSIBILITIES IN THE PEDREGOSA BASIN

By

EUGENE GREENWOOD

Midland, Texas

Oil and gas has been found in the Pedregosa Basin area. Will it be found in commercial quantities? Geologists such as Kottlowski, Wengerd, Foster, and Zeller have suggested the oil potential of various horizons. Analogies have been drawn between environmental and structural similarities of the Pedregosa basin and nearby oil-producing basins. Shows of oil have been reported from shallow water wells in southwestern New Mexico, southeastern Arizona, and the northern part of the state of Chihuahua. Zeller (1965) reported oil shows in the Big Hatchet Mountains in the Cretaceous strata and good petroliferous odor from Paleozoic rocks. The Hachita Dome well drilled in 1953 near the town of Hachita reported shows of oil from lower Paleozoic limestones. Humble Oil and Refining Company's B. A. State, drilled south of the Big Hatchet Mountains, recovered gas on a DST from Permian rocks. The Humble well was re-entered a few years ago by a group of independent oil men in an attempt to make a commercial gas well. They reported gas flow at the rate of 0.5 MMCFPD but lost the hole when attempting to acid frac the gas zone. Oil shows were reported in the Cockrell well drilled north of Coyote Hills. Sample shows have been reported in the Pemex well drilled at Los Chinos in northern Chihuahua, Mexico. Oil shows in the surface exposure of the Mississippian near Bavispe, Chihuahua, were noted by Pemex surface geologists.

If oil is found, what formations are most likely to produce?

Porosity in the middle sand member of the Cambrian Bliss Sandstone is present in the exposures at Big Hatchet Mountains (John Cys, personal commun.). Cambrian sands produce on anticlines on the eastern shelf of the Midland basin of West Texas.

Eight hundred feet of porous Silurian Fusselman Dolomite, present in the Franklin Mountains, was eroded by Middle Devonian uplift and erosion in the Big Hatchet Mountains area (Kottlowski and Pray, 1967). The eroded scarp was covered by sapropelic shale and dark cherts of the Upper Devonian. Similar conditions on the eastern shelf and central basin platform of the Permian basin of West Texas and southeastern New Mexico produce structural and stratigraphic oil.

Ordovician El Paso-Ellenburger dolomite is present and should be covered by the Devonian shale west of the Fusselman subcrop. The Ellenburger is an excellent producer on structure in the Permian basin of West Texas and southern New Mexico.

Bioherms of Mississippian crinoids occur in the Sacramento and San Andres Mountains and very massive limestone cliffs up to 500 feet thick are present in southwestern New Mexico and southeastern Arizona.

Horquilla reefs, 1200 feet thick, are described and photographed by Zeller (1965) in the Big Hatchet Mountains. The reef started growing in Desmoinesian time and grew into early Wolfcampian. Numerous reefs in West Texas

and southeastern New Mexico grew at the same time and have produced billions of barrels of oil.

Porous Epitaph Dolomite is visible in the Big Hatchet Mountains (John Cys, personal commun.), and reefs are present in the Mustang Mountains with lagoonal evaporites in the Whetstone Mountains of Cochise County, Arizona. Presence of back-reef evaporites up to 200 feet thick (Zeller, 1965) lends evidence that this Leonardian-age rock could be analogous to the prolific Abo production of southeastern New Mexico.

Lower Cretaceous beds contain rudistid, coralline, algal, bioclastic banks up to 500 feet thick in the Big Hatchet Mountains (Zeller, 1965) and 200 feet thick in the Mule Mountains of southeastern Arizona (Hayes, 1970).

A large basin with more than 25,000 feet of sedimentary rocks exists. For most of Paleozoic time, the basin had a similar history of environment and structural evolution as the Permian basin of West Texas and southeastern New Mexico. Billions of barrels of oil and tens of trillions of cubic feet of gas have been produced in the Permian basin. I believe a like amount will be found in the Pedregosa basin.

REFERENCES

- Greenwood, E., 1969, Oil and gas in the Pedregosa basin: *Oil and Gas Jour.*, v. 67, no. 40, p. 171-173.
- Hayes, P. T., 1970, Mesozoic stratigraphy of the Mule and Huachuca Mountains, Arizona: *U.S. Geol. Surv. Prof. Paper 658-A*, 27 p.
- Kottlowski, F. E., 1959, Real wildcat country—Pennsylvanian of southwest New Mexico: *Oil and Gas Jour.*, v. 57, no. 16, p. 148-151.
- _____, 1963, Paleozoic and Mesozoic strata of southwestern and south-central New Mexico: *N. Mex. Bureau Mines & Min. Res. Bull.* 79, 100p.
- _____, and Pray, L. C., 1967, Silurian outcrops of south-central and southwestern New Mexico: *Tulsa Geol. Soc. Digest*, v. 35, p. 209-230.
- _____, Foster, R. W., and Wengerd, S. A., 1969, Key oil tests and stratigraphic sections in southwest New Mexico: *N. Mex. Geol. Soc. Guidebook of the Border Region, Chihuahua and the United States*, p. 186-196.
- Wengerd, S. A., 1962, Wildcat oil prospects in southwest New Mexico: *World Oil*, v. 155, no. 2, p. 51-58.
- _____, 1969, Geologic history and exploration for oil in the border region: *N. Mex. Geol. Soc. Guidebook of the Border Region, Chihuahua and the United States*, p. 197-204.
- Zeller, R. A., Jr., 1965, Stratigraphy of the Big Hatchet Mountains area, New Mexico: *N. Mex. Bureau Mines & Min. Res. Mem.* 16, 128p.



Animas Range. View east from N.M. Highway 338 near Gray Ranch headquarters. Lack of exposures, and presence of stream terraces cut in alluvial fill, makes petroleum exploration by geological means very difficult in the Animas Valley. (Photograph by Wengerd)