



## *A model for subduction origin and distribution of fluorite deposits in the western United States*

Albert L. Lamarre

1975, pp. 169-170. <https://doi.org/10.56577/FFC-26.169>

in:

*Las Cruces Country*, Seager, W. R.; Clemons, R. E.; Callender, J. F.; [eds.], New Mexico Geological Society 26<sup>th</sup> Annual Fall Field Conference Guidebook, 376 p. <https://doi.org/10.56577/FFC-26>

---

*This is one of many related papers that were included in the 1975 NMGS Fall Field Conference Guidebook.*

---

### **Annual NMGS Fall Field Conference Guidebooks**

Every fall since 1950, the New Mexico Geological Society (NMGS) has held an annual [Fall Field Conference](#) that explores some region of New Mexico (or surrounding states). Always well attended, these conferences provide a guidebook to participants. Besides detailed road logs, the guidebooks contain many well written, edited, and peer-reviewed geoscience papers. These books have set the national standard for geologic guidebooks and are an essential geologic reference for anyone working in or around New Mexico.

### **Free Downloads**

NMGS has decided to make peer-reviewed papers from our Fall Field Conference guidebooks available for free download. This is in keeping with our mission of promoting interest, research, and cooperation regarding geology in New Mexico. However, guidebook sales represent a significant proportion of our operating budget. Therefore, only *research papers* are available for download. *Road logs*, *mini-papers*, and other selected content are available only in print for recent guidebooks.

### **Copyright Information**

Publications of the New Mexico Geological Society, printed and electronic, are protected by the copyright laws of the United States. No material from the NMGS website, or printed and electronic publications, may be reprinted or redistributed without NMGS permission. Contact us for permission to reprint portions of any of our publications.

One printed copy of any materials from the NMGS website or our print and electronic publications may be made for individual use without our permission. Teachers and students may make unlimited copies for educational use. Any other use of these materials requires explicit permission.

*This page is intentionally left blank to maintain order of facing pages.*

# A MODEL FOR SUBDUCTION ORIGIN AND DISTRIBUTION OF FLUORITE DEPOSITS IN THE WESTERN UNITED STATES

by

ALBERT L. LAMARRE  
Noranda Exploration, Inc.  
P.O. Box 15638  
Denver, Colorado 80215

## DISTRIBUTION AND AGE OF FLUORITE DEPOSITS IN THE WESTERN UNITED STATES

Fluorite occurs in each of the western states in deposits of varying sizes and importance. Figure 1, which is modified from Peters (1958), shows the distribution of fluorite deposits of hydrothermal origin in the western United States and Mexico. An obvious linear trend of deposits passes through western Montana, Wyoming, central Colorado, New Mexico, west Texas and eastern Mexico. A less clearly defined trend passes through eastern Idaho, western Utah, eastern Nevada, southern California and western Arizona.

These fluorite deposits have two common characteristics: they are middle to late Tertiary in age and are associated with alkalic igneous rocks in places where igneous rocks have been found. Specific examples of dated occurrences are given by

McAnulty and others (1963), Kottlowski (1965) and Anderson (1954). Worl (1974) gives many examples of the relationship of fluorite to alkalic igneous rocks.

## VOLCANISM AND SUBDUCTION ZONES IN THE WESTERN UNITED STATES

The middle to late Tertiary fluorite deposits are within Tertiary volcanic provinces. These provinces are characterized by rocks of intermediate calc-alkalic composition that are more alkalic toward the interior of the continent in the North and South American Cordilleran Belt (Lipman and others, 1972). The Tatil, San Juan, and Challis volcanic fields are examples of such volcanic provinces. The volcanic rocks are comprised of andesite to rhyodacite lavas commonly closely associated with younger, more silicic ash-flow sheets of latite to rhyolite composition. Similar volcanic associations are forming now at the Pacific Ocean margin, where continental plates are overriding oceanic crust in island arc environments (Hatherton and Dickinson, 1969).

Plots of  $K_2O$  versus  $SiO_2$  content for a wide variety of early and middle Tertiary igneous rock suites from the western United States (Lipman and others, 1972) fall within the family of trends defined by circum-Pacific calc-alkalic suites that are clearly related to active seismic zones and subduction of oceanic crust (Hatherton and Dickinson, 1969). This chemical similarity in  $K_2O$  versus  $SiO_2$  ratio suggests that early and middle Tertiary volcanic rocks of the western United States, like modern volcanic rocks around the Pacific basin, are genetically related to a subduction system.

Hatherton and Dickinson (1969), in studying calc-alkalic rocks overlying subduction zones in the Lesser Antilles, New Zealand, and Indonesia found that for volcanic rocks with 60 percent silica the  $K_2O$  content of the rocks increased linearly with increasing depth to the underlying subduction zone. Lipman and others (1972) used this  $K_2O$ -depth relationship by determining  $K_2O$  content at 60 percent  $SiO_2$  for 70 known early and middle Tertiary igneous rock suites from the western United States.  $K_2O$  contents were converted to depth values and plotted on a map of the western United States, and then contoured at 50 km intervals. The plot (Fig. 2) shows two well-defined trends of eastward increasing depths to subduction zones and suggests that early and middle Tertiary subduction of the Pacific Ocean crust beneath western North America was along two imbricate subduction zones dipping eastward at  $15^\circ$  to  $20^\circ$ .

The calc-alkalic igneous rocks, and the more alkalic products, are postulated to originate by melting of the oceanic lithospheric slab along subduction zones beneath continental crust. Upon subduction of the slab, dehydration reactions take place in response to increasing temperature and pressure, and various metamorphic minerals become unstable and dissociate




Figure 1. Distribution of fluorite mines and occurrences in the western United States (modified from Peters, 1958). Special attention is drawn to the dots, representing fluorite mines, which exemplify the linear trends of fluorite distribution. The x's represent occurrences in which fluorite may be only a small component.




Figure 2. Contoured depths (km) to inferred early and middle Tertiary subduction zones of the western United States. Double line indicates discontinuity in contours. From Lipman and others, 1972.

at depth. Dissociated serpentine, talc, biotite, muscovite and hornblende may react with anhydrous mantle peridotite and water to produce phlogopite (Brown and Fyfe, 1972). Phlogopite is a relatively K-rich mineral which may be stable to a depth of about 300 km (Wyllie, 1973). Melting at depths greater than 300 km would therefore be expected to produce K-rich igneous rocks and may explain the increasing alkalinity of subduction-related igneous rocks continentward from oceanic trenches (Wyllie, 1971). Furthermore, phlogopite may contain up to 6.74 percent fluorine (Allman and Koritnig, 1972).

#### RELATIONSHIP OF FLUORITE DEPOSITS TO SUBDUCTION ZONES

Comparison of Figures 1 and 2 shows the distribution of middle to late Tertiary fluorite deposits is nearly coincident with the surface projection of the easternmost part of each of the two early to middle Tertiary subduction zones as described by Lipman and others (1972). I suggest that the fluorine in hydrothermal fluorite deposits in the western United States is

derived by melting of phlogopite along the deeper parts of Tertiary subduction zones. Because of the ability of alkalic melts to hold fluorine as alkali-fluoride compounds (Kogarko and others, 1968) the melts tend to concentrate fluorine along subduction zones and transport it upward through the mantle. In the crust fluorine volatilizes as  $\text{SiF}_4$  (Kogarko and others, 1968) and may pass upward through tensional faults, where it mixes with meteoric water to form  $\text{HF}$ - and  $\text{SiO}_2$ -bearing hydrothermal fluids. Fluorite may precipitate from these fluids if physiochemical and structural conditions are favorable.

Fluorite deposits in the western United States are therefore found within or in close proximity to tensional fault zones which overlie the positions of the deepest parts of the two early to middle Tertiary subduction zones. The Rio Grande rift is a good example of an area of calc-alkalic volcanism and tensional faulting having abundant fluorite deposits.

#### REFERENCES

- Allman, R., and Koritnig, S., 1972, Fluorine, Ch. 9: in Wedepohl, K. H., ed., *Handbook of Geochemistry* 11-3.
- Anderson, A. L., 1954, A preliminary report on the fluorspar mineralization near Challis, Custer County, Idaho: Idaho Bur. Mines and Geol. Pamphlet 101, 13 p.
- Brown, G. C., and Fyfe, W. S., 1972, The transition from metamorphism to melting: status of the granulite and eclogite facies: 24th I.G.S., Section 2, p. 27-34.
- Hatherton, T., and Dickinson, W. R., 1969, The relationship between andesitic volcanism and seismicity in Indonesia, the Lesser Antilles and other island arcs: *Jour. Geophys. Res.*, v. 74, p. 5301-5310.
- Kogarko, L. N., Krigman, L. D., and Sharudilo, N. S., 1968, Experimental investigations of the effect of alkalinity of silicate melts on the separation of fluorine into the gas phase: *Geochem. International*, v. 5, p. 782-790.
- Kottlowski, F. E., 1965, Middle Oligocene fluorite-barite mineralization in New Mexico (abs.): *Geol. Soc. America Annual Meeting, Program with Abstracts*, p. 90.
- Lipman, P. W., Prostka, H. J., and Christiansen, R. L., 1972, Cenozoic volcanism and plate-tectonic evolution of the western United States, part I. early and middle Cenozoic: *Phil. Trans. R. Soc. London*, v. 271, p. 217-248.
- McAnulty, W. N., Sewell, C. R., Atkinson, D. R., and Rasberry, J. M., 1963, Aguachile beryllium-bearing fluorspar district, Coahuila, Mexico: *Geol. Soc. America Bull.*, v. 74, p. 735-744.
- Peters, W. C., 1958, Geologic characteristics of fluorspar deposits in the western United States: *Econ. Geol.*, v. 53, p. 663-688.
- Worl, R. G., 1974, Geology of fluorspar deposits of the western United States, *in* A Symposium on the Geology of Fluorspar: *Kentucky Geol. Surv. Spec. Pub.* 22, p. 31-54.
- Wyllie, P. J., 1971, *The Dynamic Earth*: Toronto, John Wiley & Sons, Inc., 416 p.
- , 1973, Experimental petrology and global tectonics—a preview: *Tectonophysics*, v. 17, p. 189-209.