



## *Tectonics of thrust and fold belt of northwestern Chihuahua*

L. L. Corbitt

1988, pp. 67-70. <https://doi.org/10.56577/FFC-39.67>

in:

*Cretaceous and Laramide Tectonic Evolution of Southwestern New Mexico*, Mack, G. H.; Lawton, T. F.; Lucas, S. G.; [eds.], New Mexico Geological Society 39<sup>th</sup> Annual Fall Field Conference Guidebook, 216 p.

<https://doi.org/10.56577/FFC-39>

---

*This is one of many related papers that were included in the 1988 NMGS Fall Field Conference Guidebook.*

---

### **Annual NMGS Fall Field Conference Guidebooks**

Every fall since 1950, the New Mexico Geological Society (NMGS) has held an annual [Fall Field Conference](#) that explores some region of New Mexico (or surrounding states). Always well attended, these conferences provide a guidebook to participants. Besides detailed road logs, the guidebooks contain many well written, edited, and peer-reviewed geoscience papers. These books have set the national standard for geologic guidebooks and are an essential geologic reference for anyone working in or around New Mexico.

### **Free Downloads**

NMGS has decided to make peer-reviewed papers from our Fall Field Conference guidebooks available for free download. This is in keeping with our mission of promoting interest, research, and cooperation regarding geology in New Mexico. However, guidebook sales represent a significant proportion of our operating budget. Therefore, only *research papers* are available for download. *Road logs*, *mini-papers*, and other selected content are available only in print for recent guidebooks.

### **Copyright Information**

Publications of the New Mexico Geological Society, printed and electronic, are protected by the copyright laws of the United States. No material from the NMGS website, or printed and electronic publications, may be reprinted or redistributed without NMGS permission. Contact us for permission to reprint portions of any of our publications.

One printed copy of any materials from the NMGS website or our print and electronic publications may be made for individual use without our permission. Teachers and students may make unlimited copies for educational use. Any other use of these materials requires explicit permission.

*This page is intentionally left blank to maintain order of facing pages.*

# TECTONICS OF THRUST AND FOLD BELT OF NORTHWESTERN CHIHUAHUA

L. L. CORBITT

Consulting Geologist, Midland, Texas 79701

**Abstract**—The Cordilleran thrust and fold belt of northwestern Chihuahua is characterized by flat-lying thrusts and closely compressed folds. These structures are of Laramide age (late Cretaceous-early Tertiary) and are well exposed in northwestern Chihuahua in the Sierras Rica, Palomas (Alta), Boca Grande, Chinos, Santa Rita and Salada. Rocks deposited prior to and during Laramide deformation are exposed as small patches surrounded by younger post-orogenic volcanic rocks of mid-Tertiary age and late Cenozoic clastic sediments and basalt. Yielding on the thrust faults and folds is northeastward toward the foreland with displacements as much as several km. Basin and Range extensional block faulting occurred in late Cenozoic time, and range marginal faults truncate some of the Laramide structures.

## INTRODUCTION

The thrust and fold belt of northwestern Chihuahua (Alvarez, 1949; Fig. 1) is well exposed in the Sierras Rica, Palomas, Boca Grande, Chinos, Santa Rita, Salada, Monumento, Cartucho, Borregos, Barrial and Nariz. Major overthrusts are exposed in the Sierras Rica and Salada. The other ranges are characterized by flat-lying thrusts of lesser displacement and closely compressed folds. Surface, subsurface, seismic (Harvey Sales) and magnetic data indicate that Paleozoic rocks have been thrust to the northeast over Cretaceous strata in the northern Animas Mountains, the Big Hatchet Mountains and Sierra Rica in the U.S., and in the Sierras Rica and Salada, in Mexico, along the Sierra Rica overthrust.

Drewes (1982) includes the area between the southern Florida Mountains (Fig. 1) and the Tres Hermanas Mountains in southwestern New Mexico in his para-autochthonous zone of minor thrusting. Brown and Clemons (1983) and Seager and Mack (1986) interpret the chief mode of Laramide deformation in southwestern New Mexico to be uplift of

relatively simple basement blocks and subsidence of complementary basins.

S. Thompson, III (written commun. 1984) pointed out that seismic data (Harvey Sales) and well data show a 1500 m (5000 ft) deep Paleocene-Eocene molasse trough trending southeastward north of the Victorio Mountains, Snake Hills, Florida Mountains and East Potrillo Mountains.

## STRATIGRAPHY

Precambrian basement does not crop out in northwestern Chihuahua but was penetrated in two Pemex wells. Granite gneiss in the Chinos No.1 was dated by the rubidium-strontium method as  $1327 \pm 242$  my and as  $890 \pm 32$  my in the Moyotes No. 1 (Thompson et al., 1978).

Paleozoic strata are about 4270 m (14,000 ft) thick and consist mostly of platform limestone and dolomite (Fig. 2, thicknesses from Zeller, 1965). Mississippian through Permian strata are exposed in the Sierra Palomas (Sivils and Phillips, 1986). Lower Paleozoic rocks crop out




FIGURE 1. Generalized tectonic map of northwestern Chihuahua and southwestern New Mexico and west Texas, modified from Woodward and DuChene (1982).

Cenozoic	Tertiary	Clastic basin-fill		Ts		
		Rhyolitic-lafitic volcanics		Tv		
		Hidalgo Volcanics	5,500 ft	Th		
		Ringbone Formation	7,500 ft	TKr		
Mesozoic	Cretaceous	Mojado Formation		Km		
		U-Bar Formation		Ku		
		Hell-to-Finish Formation		Kh		
		Hell-to-Finish Formation		6,000 ft		
Paleozoic	Permian	Hueco Group	Concha Limestone	1,375 ft	Pc	
			Scherrer Formation	20 ft	Ps	
			Epitaph Dolomite	1,500 ft	Pe	
			Collina Limestone	500 ft	Pcl	
			Earp Formation	1,000 ft	Per	
			Horquilla Limestone	3,500 ft	Ph	
	Pennsylvanian	Paradise Formation		320 ft	Mp	
	Mississippian	Escabrosa Group	1,260 ft		Hatchita Formation	Mh
			Keating Formation			
	Devonian	Percha Shale		280 ft	Dp	
	Silurian	Fusselman Dolomite		900 ft	Sf	
	Ordovician	Montoya Formation		385 ft	Om	
		El Paso Formation		1,070 ft	Oep	
	Cambrian	Bliss Formation		327 ft	O C b	
	Precambrian	Granitic and Metamorphic Rocks			p C	

FIGURE 2. Composite stratigraphic sequence in southwestern New Mexico and northwestern Chihuahua, Mexico, adapted from Woodward and DuChene (1982).

in the Sierras Rica, Salada and Monumento. Permian rocks crop out in the Sierras Chinos and Santa Rita.

Triassic rocks are absent in northwestern Chihuahua. Jurassic rocks are reported in the Pemex Moyotes and Sapallo tests (Fig. 1), and are mapped in the Sierra Aguila south of Palomas (Reyeros de Castillo, 1974; Rodriguez, 1975). The unnamed evaporites underlying the Hell-to-Finish Formation in the Little Hatched Mountains (Zeller, 1970) and the intrusive gypsum pods reported by Drewes (1982) in the Animas Mountains, also may be Jurassic in age.

Lower Cretaceous strata in northwestern Chihuahua have a maximum thickness of about 3050 m (10,000 ft) and include clastic red beds (Hell-to-Finish) and limestone with rudistids (U-Bar). These formations are best exposed in the Sierras Boca Grande and Cartucho.

Lower Tertiary rocks are of volcanic and continental clastic origin and probably accumulated contemporaneously with development of Laramide thrusts and folds. Mid-Tertiary (Oligocene and younger) volcanic rocks are widespread. Coarse clastic sediments of late Cenozoic age fill the basins between the present mountain ranges. Late Cenozoic basaltic rocks are locally present (Frantes, 1981).

### PALEOTECTONIC SETTING

From Cambrian through Mississippian time, northwestern Chihuahua was part of a stable shelf south of the transcontinental arch. Three episodes of epeirogenic uplift, during the Middle Ordovician, Early Silurian and Late Silurian-Middle Devonian, are recorded by stratigraphic breaks. In Pennsylvanian time, the area was epeirogenically deformed with development of the Alamo Hueco basin (Zeller, 1965) in which 610 m (2,000 ft) or more of basinal sediments were deposited. The basin was bordered on the northeast by the Florida-Moyotes-Diablo uplift. Permian Hueco Formation overlies Mississippian limestone in the Florida Mountains. Abo sediments overlie Precambrian rocks in the Pemex Moyotes test and Permian rocks were deposited on Precambrian and Paleozoic rocks on the Diablo uplift. Paleozoic strata appear to have been removed from the Burro uplift in Mesozoic time (Elston, 1958). Pennsylvanian-Permian basinal sediments are recognized in the Humble No. 1 State BA (sec. 25, T32S, R16W) and in the Pemex Villa Ahumada test. Further subsidence of the Pedregosa basin occurred during the Permian with accumulation of over 1830 m (6,000 ft) of shelf deposits along the margin of the basin.

Turner (1962) described the Deming axis (Florence-Graham-Burro-Florida-Diablo uplift of Greenwood et al., 1977) as being a positive element throughout much of Mesozoic time. Cretaceous sandstone overlies Precambrian rocks in the Burro Mountains, Fusselman Dolomite in the Victorio Mountains and in the Marshall Young Bisbee Hills test south of the Snake Hills (Fig. 1; S. Thompson, III, written commun. 1986), Mississippian rocks in the Mobil test south of El Paso and Precambrian and Paleozoic rocks on the Diablo uplift. The Late Jurassic and Early Cretaceous Chihuahua trough was present south of the Deming Axis (DeFord and Haenggi, 1970, fig. 5).

### TECTONIC FRAMEWORK

The overthrust mapped by Zeller (1958) in the Sierra Rica is exposed to the southeast in Mexico (Fig. 1). The view of the southwest-dipping, dark-colored, lower Paleozoic rocks overlying the lighter Cretaceous strata is striking. Approximately eight km southeast of the Sierra Rica in the El Gato hills, Silurian Fusselman rocks appear to be thrust over Mississippian carbonates. These outcrops dip to the southwest and are 10 km west of Permo-Pennsylvanian rocks exposed in the Sierra Palomas. These rocks also dip to the southwest. It is possible that the strata exposed in the El Gato hills are thrust over the Permo-Pennsylvanian rocks exposed in the Sierra Palomas.

The Pemex Camello test, 16 km to the south of El Gato (Fig. 1), drilled in repeated Cretaceous rocks at a total depth of 6000 m (18,352 ft). At least two thrusts are reported in this test (Thompson et al., 1978). This well is located about 16 km east of the Big Hatchet Mountains where Permo-Pennsylvanian carbonates are more than 2440 m (8,000 ft) above sea level. Precambrian basement in the Camello test would appear to be at a depth of approximately 10,000 m (33,000 ft) based on the 4267 m (14,000 ft) of Paleozoic rocks drilled in the Humble No. 1 State BA test, 29 km to the west, and in the Pemex Chinos well, 32 km to the southeast (Fig. 1).

The Big Hatchet Mountains and Sierra Rica appear to be thrust over Cretaceous strata along the Sierra Rica overthrust. The Pemex Camello test is located immediately east of this thrust and drilled Cretaceous strata similar to that exposed in the Little Hatchet Mountains and the northern Sierra Rica. The Big Hatchet Mountains overlie a regional down-to-the-southwest magnetic gradient with no indication of magnetic basement involvement. Seismic data (Harvey Sales) along the border between the Big Hatchet Mountains and Pemex Camello test do not indicate any major normal faulting. Also, a thrust appears to approach the surface in the Sierra Rica on this seismic data.

Seager (1983) suggests that the Big Hatchet Mountains are a Rocky Mountain type upthrust with Laramide basins present to the north and south. Synorogenic rocks are present to the north in the Little Hatchet Mountains but not to the south. The Cockrell tests drilled west of the Big Hatchet Mountains indicate this area was more deeply eroded than the Big Hatchet Mountains before Late Tertiary Basin and Range faulting (Fig. 3; Woodward and DuChene, 1982). Seismic (Harvey Sales), magnetic and gravity data indicate that the Basin and Range normal fault on the west side of the Big Hatchet Mountains flattens with depth and that Precambrian basement may not be involved.

The leading edge of the Sierra Rica overthrust appears to lie between the Big Hatchet Mountains and the Pemex Camello well and may be exposed to the south in the Sierra Salada. A structurally complex thrust plate of Paleozoic strata dipping to the southwest overlies Lower Cretaceous rocks in the Sierra Salada.

Northwest of the Sierra Salada at least three exposures of lower Paleozoic rocks are present in the Sierra Monumento. These strata are surrounded by mid-Tertiary volcanics and are folded into northwest-trending folds characteristic of northwestern Chihuahua. Complex structural relations are also indicated in the Centuro test drilled west of Sierra Salada, which drilled 3294 m (10,800 ft) of Permian rocks (Thompson et al., 1978).

A small exposure of tightly folded Cretaceous strata is present about 24 km east of Sierra Salada near the Pemex Ascension test (Fig. 1). Very complex overturned folds and thrusts in upper Paleozoic rocks are present at the north end of the Sierra Palomas. Broad northwest-trending folds dominate the rest of the range. Complex overturned folds




FIGURE 3. Diagrammatic cross section of the thrust belt of southwestern New Mexico. See Figure 1 for line of section.

in upper Paleozoic rocks are also present to the south in the valley between Sierra Palomas and Boca Grande, indicating yielding to the northeast. Boca Grande is a syncline similar to the smaller U-Bar syncline south of the Big Hatchet Mountains. Very tight folds dominate the Sierra Chinos (Brown and Dyer, 1986). Numerous small imbricate thrust faults and tight folds are present in the Sierra Santa Rita. This range of Permian rocks may be thrust to the northeast over Cretaceous strata. Several major folds and thrusts are exposed in Sierra Cartucho and to the southeast in the Sierras Borregos, Barrial and Nariz.

### CONCLUSIONS

The evidence suggests that northwestern Chihuahua is underlain by a regional system of thrust faults and folds that have yielded toward the northeast. Total displacement on the Sierra Rica overthrust is not known but is probably several km.

The Marshall Young No. 1 Salty Unit well (sec. 33, T23S, R15W), Grant County, New Mexico (Fig. 3) appears to verify the presence of the overthrust belt in southwestern New Mexico (Brennan, 1986). The following section was drilled in this test: surface Tertiary volcanics: 216 m (710 ft); Tertiary conglomerate: 235 m (770 ft); Epitaph Formation (Permian): 311 m (1020 ft); Colina Limestone: 356 m (1168 ft); Earp Formation: 657 m (2154 ft); Horquilla Formation (Pennsylvanian): 811 m (2660 ft); Escabrosa Formation: 1084 m (3558 ft); Percha Shale (Devonian): 1144 m (3756 ft); Montoya Formation (Ordovician): 1292 m (4238 ft); El Paso Formation: 1361 m (4466 ft); Bliss Formation (Cambrian-Ordovician): 1369 m (4494 ft); thrust fault; Mojado Formation (Lower Cretaceous): 2306 m (7564 ft); U-Bar Formation: 2348 m (7705 ft); Hell-to-Finish Formation: 2543 m (8344 ft); unconformity; Montoya Formation: 2713 m (8960 ft); El Paso Formation: 2962 m (9718 ft); Bliss Formation to T.D. (S. Thompson, III,

written commun. 1986). Several kilometers of tectonic transport to the northeast is suggested.

To the southeast in the Tres Hermanas Mountains and East Potrillo Mountains Permian rocks have been thrust to the northeast over Cretaceous rocks. The following section was drilled in the Pure test (Fig. 1) east of the East Potrillo Mountains (Thompson, 1982): surface; Cretaceous: 190 m (625 ft); Permian? to Mississippian?: 1174 m (3850 ft); Percha: 1238 m (4060 ft); Fusselman: 1343 m (4405 ft); reverse fault?; Permian? to Mississippian? rocks: 1903 m (6240 ft); marble?: 2078 m (6815 ft); intrusive rocks. Several kilometers of northeast transport is indicated.

In the Juarez Mountains (Fig. 1) Lovejoy (1980) estimated a minimum northeast transport of 22 km (13.2 mi) and that the original width of Cretaceous strata had been decreased 59% by thrusting and folding.

In the Malone Mountains in Texas, Berg (1982) estimated a combined northeast tectonic transport of at least 24 km (15 mi) and that it may have been as much as 80 km (50 mi). In the Quitman Mountains and the Cieneguilla Mountains in Mexico, Reaser (1982) estimated 48 km (30 mi) of northeast tectonic transport. Similar amounts of northeast tectonic transport are probably present in northwestern Chihuahua.

The Laramide thrust and fold belt appears to have developed to the northeast of a continental-margin volcanic arc (Lipman and Sawyer, 1985). Mack (1987) indicates that a retroarc foreland basin formed in northwestern Chihuahua-southwestern New Mexico in late Albian-Cenomanian time, northeast of a complementary compressional orogenic belt. Possibly a gravitational gliding model similar to that proposed by Price and Mountjoy (1970) may have been operative with upward and outward pushing from the volcanic arc. The Laramide volcanic arc area was later covered by mid-Tertiary volcanics which make up the present-day Sierra Madre Occidental (Damon et al., 1981).

## ACKNOWLEDGMENTS

I thank R. E. Clemons, L. A. Woodward and Russ Dyer for critically reviewing the manuscript.

## REFERENCES

- Alvarez, M., 1949, Tectonics of Mexico: American Association of Petroleum Geologists Bulletin, v. 33, p. 1484-1469.
- Berg, T. P., 1982, Structural evaluation of the northeastern Chihuahua tectonic belt; *in* Powers, R. B., ed., Geologic studies of the Cordilleran thrust belt: Rocky Mountain Association of Geologists Guidebook, p. 451-485.
- Brennan, D. J., 1986, Oil and gas developments in Four Corner-Intermountain area in 1983-1985: American Association of Petroleum Geologists Bulletin, v. 70, p. 1315-1319.
- Brown, C. A. and Clemons, R. E., 1983, Florida Mountains section of southwest New Mexico overthrust belt—a reevaluation: New Mexico Geology, v. 5, p. 26-29.
- Brown, M. L. and Dyer, R., 1986, Structural geology of Sierra de los Chinos and Cerro la Cueva, northwestern Chihuahua, Mexico; *in* Hoffer, J. M., ed., Geology of south-central New Mexico: El Paso Geological Society Guidebook, p. 141-151.
- Damon, P. E., Shafiqulla, M. and Clark, K., 1981, Age trends of igneous activity in relation to metallogenesis in the southern Cordillera: Arizona Geological Society Digest, v. 14, p. 137-154.
- DeFord, R. and Haeggi, W. T., 1970, Stratigraphic nomenclature of Cretaceous rocks in northeastern Chihuahua; *in* The geologic framework of the Chihuahua tectonic belt: West Texas Geological Society, p. 175-196.
- Drewes, H., 1982, Commentary and roadlog of field trip El Paso, Texas to Wickenburg, Arizona; *in* Powers, P. B., ed., Geologic studies of Cordilleran thrust belt: Rocky Mountain Association of Geologists Guidebook, v. 2, p. 895-930.
- Drewes, H., 1982, Some general features of the El Paso-Wickenburg transect of the Cordilleran orogenic belt, Texas to Arizona; *in* Powers, P. B., ed., Geological studies of Cordilleran thrust belt: Rocky Mountain Association of Geologists Guidebook, v. 2, p. 882-894.
- Elston, W. E., 1958, Burro uplift, northeastern limit of sedimentary basin of southwestern New Mexico and southeastern Arizona: American Association of Petroleum Geologists Bulletin, v. 42, p. 2513-2517.
- Frantes, T. J., 1981, The geology of the Palomas volcanic field, Luna County, New Mexico and Chihuahua [M.S. thesis]: El Paso, University of Texas at El Paso, 70 p.
- Greenwood, E., Kottowski, F. E. and Thompson, S., III, 1977, Petroleum potential and stratigraphy of Pedregosa basin: Comparison with Permian and Orogrande basins: American Association of Petroleum Geologists Bulletin, v. 61, 1448-1469.
- Lipman, P. W. and Sawyer, D. A., 1985, Mesozoic ash-flow caldera fragments in southeastern Arizona and their relation to porphyry copper deposits: Geology, v. 13, p. 652-656.
- Lovejoy, E. M. P., ed., 1980, Sierra de Juarez, Chihuahua Mexico, structure and stratigraphy: El Paso Geological Society, 59 p.
- Mack, G. H., 1987, Mid-Cretaceous (late Albian) change from rift to retroarc foreland basin in southwestern New Mexico: Geological Society of America Bulletin, v. 98, p. 507-514.
- Price, R. A. and Mountjoy, E. W., 1970, Geologic structure of the Canadian Rocky Mountains between Bow and Athabasca Rivers—a progress report; *in* Structure of the southern Canadian Cordillera: Geologic Association Canada, Special Paper 6, p. 966-992.
- Reaser, D. F., 1982, Geometry and deformational environment of the Cienguilla-Quitman range in northeastern Chihuahua, Mexico, and Western Trans-Pecos, Texas, USA; *in* Powers, R. B., ed., Geologic studies of the Cordilleran thrust belt: Rocky Mountain Association of Geologists Guidebook, p. 425-449.
- Reyerros de Castillo, M. M., 1974, Corales del Jurassic Superior de Chihuahua: Paleontologia Mexicana, No. 40, Institute de Geologia U.N.A.M., p. 7-43.
- Rodriguez, J. T., 1975, Preliminary geologic map of the northeastern section of the state of Chihuahua; *in* Exploration from the mountains to the basin: El Paso Geologic Society.
- Seager, W. R., 1983, Laramide wrench faults, basement-cored uplifts, and complementary basins in southern New Mexico: New Mexico Geology, v. 5, p. 69-76.
- Seager, W. R. and Mack, G. H., 1986, Laramide paleotectonics of southern New Mexico: American Association of Petroleum Geologists, Memoir 41, p. 669-685.
- Sivils, D. J. and Phillips, J. D., 1986, Geology of Sierra de Palomas, Chihuahua Mexico; *in* Hoffer, J. M., ed., Geology of south-central New Mexico: El Paso Geologic Society Guidebook, p. 60-66.
- Thompson, S., III, 1982, Oil and gas exploration wells in southwestern New Mexico; *in* Powers, R. P., ed., Geologic studies of the Cordilleran thrust belt: Rocky Mountain Association of Geologists, p. 521-536.
- Thompson, S., III, Tovar, J. and Conley, J. N., 1978, Oil and gas exploration wells in the Pedregosa basin: New Mexico Geological Society, Guidebook 29, p. 331-342.
- Turner, G. L., 1962, The Deming axis of southeastern Arizona, New Mexico, and Trans-Pecos Texas: New Mexico Geological Society, Guidebook 13, p. 59-71.
- Woodward, L. A. and DuChene, H. R., 1982, Tectonics and hydrocarbon potential of thrust and fold belt, southwestern New Mexico; *in* Powers, R. P., ed., Geologic studies of the Cordilleran thrust belt: Rocky Mountain Association of Geologists Guidebook, v. 1, p. 409-418.
- Zeller, R. A., 1958, The geology of Big Hatchet Peak quadrangle, Hidalgo County, New Mexico [Ph.D. dissertation]: Los Angeles, University of California, 260 p.
- Zeller, R. A., 1965, Stratigraphy of the Big Hatchet Mountain area New Mexico: New Mexico Bureau of Mines and Mineral Resources, Memoir 16, 128 p.
- Zeller, R. A., 1970, Geology of the Little Hatchet Mountains, Hidalgo and Grant Counties, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Bulletin 96, 23 p.