



Preliminary observations on the mining history, geology, mineralization of the Jicarilla mining district, Lincoln County, New Mexico

Virginia T. McLemore, Mark Ouimette, and Robert W. Eveleth
1991, pp. 311-316. <https://doi.org/10.56577/FFC-42.311>

in:
Geology of the Sierra Blanca, Sacramento, and Capitan Ranges, New Mexico, Barker, J. M.; Kues, B. S.; Austin, G. S.; Lucas, S. G.; [eds.], New Mexico Geological Society 42nd Annual Fall Field Conference Guidebook, 361 p.
<https://doi.org/10.56577/FFC-42>

This is one of many related papers that were included in the 1991 NMGS Fall Field Conference Guidebook.

Annual NMGS Fall Field Conference Guidebooks

Every fall since 1950, the New Mexico Geological Society (NMGS) has held an annual [Fall Field Conference](#) that explores some region of New Mexico (or surrounding states). Always well attended, these conferences provide a guidebook to participants. Besides detailed road logs, the guidebooks contain many well written, edited, and peer-reviewed geoscience papers. These books have set the national standard for geologic guidebooks and are an essential geologic reference for anyone working in or around New Mexico.

Free Downloads

NMGS has decided to make peer-reviewed papers from our Fall Field Conference guidebooks available for free download. This is in keeping with our mission of promoting interest, research, and cooperation regarding geology in New Mexico. However, guidebook sales represent a significant proportion of our operating budget. Therefore, only *research papers* are available for download. *Road logs*, *mini-papers*, and other selected content are available only in print for recent guidebooks.

Copyright Information

Publications of the New Mexico Geological Society, printed and electronic, are protected by the copyright laws of the United States. No material from the NMGS website, or printed and electronic publications, may be reprinted or redistributed without NMGS permission. Contact us for permission to reprint portions of any of our publications.

One printed copy of any materials from the NMGS website or our print and electronic publications may be made for individual use without our permission. Teachers and students may make unlimited copies for educational use. Any other use of these materials requires explicit permission.

This page is intentionally left blank to maintain order of facing pages.

PRELIMINARY OBSERVATIONS ON THE MINING HISTORY, GEOLOGY AND MINERALIZATION OF THE JICARILLA MINING DISTRICT, LINCOLN COUNTY, NEW MEXICO

VIRGINIA T. McLEMORE^{1,2}, MARK OUIMETTE² and ROBERT W. EVELETH¹

¹New Mexico Bureau of Mines and Mineral Resources, Socorro, New Mexico 87801; ²Department of Geological Sciences, University of Texas at El Paso, El Paso, Texas 79968

Abstract—The Jicarilla mining district comprises much of the Jicarilla Mountains in central New Mexico. The three types of mineral deposits that occur in the district are placer, quartz veins and iron skarn/replacement deposits. Production has been minor and amounts to less than 800 oz of placer gold, 800 oz of lode gold, 38,000 oz of silver, 4.2 million lbs of copper, and 3000 lbs of lead. In addition, about 8000 tons of iron ore were produced in the 1900s. Mining began two or three hundred years ago and the district was perhaps the site of the first drill-rig in New Mexico in 1881. The mineral deposits appear to be related to Tertiary intrusives that are diverse in lithology and composition. Two age dates of about 37–38 Ma were obtained on samples of the igneous rocks. The origin of the mineral deposits was a complex process of magmatic fractionation and differentiation, perhaps involving upper mantle and lower crustal sources. The placer deposits are large but low grade, and the scarcity of water prevents developments. The district has potential for gold-bearing veins and possible breccia deposits, but subsurface drilling will be required to determine their extent.

INTRODUCTION

The Jicarilla (or Ancho) mining district is located about 18 km north-east of White Oaks in northern Lincoln County, central New Mexico (Fig. 1). The district comprises much of the Jicarilla Mountains. The geology and mineralization have been described in several reports (Jones, 1904; Lindgren et al., 1910; Wright, 1932; Griswold, 1959; Budding, 1963, 1964; Ryberg, 1968; Johnson, 1972; Segerstrom and Ryberg, 1974; North and McLemore, 1986). However, very little published information is available regarding the history of this mining district, which was one of the earliest active districts in New Mexico. The purpose of this paper is threefold: (1) to discuss the mining history of the district; (2) to briefly describe the geology, igneous petrology, and different types of mineral deposits; and (3) to relate the petrogenesis and mineralization to similar types of deposits in New Mexico. This work is based upon a literature search, field reconnaissance and preliminary petrologic and geochemical studies. These data are incomplete and the results are preliminary.

MINING HISTORY AND PRODUCTION

Although placer mining in the Jicarilla mining district has been carried on at least since the mid-nineteenth century, total known production is disappointingly low. Some accounts indicate the placers were first worked during the Spanish occupation, two to three hundred years ago (Engineering and Mining Journal, 5/19/1904, p. 799; Bullion, 7/26/1892, p. 4). Jones (1904, p. 177) suggested that native Mexican miners were working the ground as early as 1850. Such mining was difficult at best due to the scarcity of water, and was carried on in a most primitive manner. Gold-bearing material was carried some distance to where water was available (occasionally it, too, being transported with great effort) and the material washed in "bateas," the predecessor of the gold pan. So extensive was this activity that by the time of the arrival of American prospectors around 1880, Ancho and Rico gulches were said to be honeycombed with placer diggings for 10 km (Mining World, vol. 1, no. 10, June 1881, p. 9; Jones, 1904, p. 177).

Overall grade of the placers was low but local "hot" spots were apparently found. Miguel Antonio Otero (father of New Mexico governor Miguel Otero) and his associates are said to have worked a crew of miners during the 1860 placering season and recovered some 2900 oz. worth approximately \$60,000 (Bullion, 7/26/92, p. 4). The local high-grade streak could not possibly account for all this activity, however. Jicarilla gold is exceptionally fine and brought a dollar more per ounce than most other placer gold in the territory—a significant fact whether the price of the metal is \$20.67 per oz (as it was in the nineteenth and early twentieth centuries) or \$400 per oz. Even the slightest amount of precipitation, whether rain or snow, would bring the miners out in force to wash for the yellow metal (Mining World, June 1881, p. 9).

The first attempts at lode mining began with the arrival of the American prospectors. Some of the earliest properties located were the Argonaut, Buckeye State, Eureka, Gold Stain (now Spring), Hawkeye, Red Rose and Zulu. Also during this period, New Mexico's very first drill-rig may have been set up. According to Mining World (Las Vegas, NM), a large boring machine was shipped from the oil regions of Pennsylvania to Las Vegas by rail and thence by teams to Jicarilla. A pipeline was constructed to Patos Mountain to obtain enough water to run the steam-driven drill and 500 cords of wood were delivered to the site for fuel. By June 1881 the derrick was erected. Some \$15,000 was invested by the adventurers, but it is not known whether any drilling was actually completed (Mining World, March 1881, p. 12; June 1881, p. 8).

The most significant developments in the district came soon after the turn of the century. The American Placer Company erected a placering machine and attempted to work the gravels principally along Ancho Gulch. Contrary to popular belief, this machine was not a dredge in the strictest sense since it did not float in water. Research shows it to have been a cantilever-type placering machine manufactured by W. M. Johnston of Chicago. The machine weighed nearly 200 tons and supposedly could excavate down to 8 m. It was a brute to move, being rail-mounted on two parallel track beds. Lack of water was again a major problem, and to obtain a sufficient supply, some 6 km of pipeline was installed by the company. Wells were sunk to 980–1320 m, but regardless of the effort and investment (said to be at least \$50,000), the scarcity of water and the difficulty of working the ground (2 to 7 m of overburden lay above the narrow pay streak and the machine obviously encountered difficulty here), the venture was unsuccessful (Jones, 1904, p. 177; Lindgren et al., 1910, p. 184; Griswold, 1959, p. 79; Engineering and Mining Journal, 9/01/01, p. 259, 11/03/01, p. 530).

Attempts were again made about the same time to develop the lode deposits. The Wisconsin Mining and Milling Company was perhaps the most successful in that some significant tonnages were developed and some bullion produced. What is not known is whether these efforts were profitable. Wisconsin installed a 50 ton/day Fairbanks-Morse amalgamation mill and focused their efforts on such properties as the Sally, Murphy, Apex and Gold Stain. Extensive development was carried on at the Gold Stain and Murphy, and today the largest mine dumps in the district are located on these properties (Western Mining World, 7/01/05, p. 701; Griswold, 1959, p. 79). Wisconsin's best years appear to have occurred during 1908–1909, when the Murphy was developed down to at least the 400-ft level and substantial tonnages of gold-bearing ores were treated (Western Mining World, 9/12/08, p. 420; Henderson, 1908, 1910).

Other attempts at lode and placer mining were made during this




FIGURE 1. Geologic map of Jicarilla mining district (simplified from Segerstrom and Ryberg, 1974).

period by such companies as the Revenue Gold Mining Co. (appropriately misnamed), Rincon Mining and Milling Co., Rico Mining Co., Fleming-Fox Copper Co., Mitchell Gold Co., and others (NMBM&MR file data; *Western Mining World*, 6/24/05, p. 673; 7/01/06, p. 701; 9/12/08, p. 420). All appear to have been marginally successful at best. The last lode mining took place in the Jicarilla district during 1933, when the Lucky Strike and a few other small prospects shipped 130 tons of gold ore (Anderson, 1957, p. 91).

The devaluation of the dollar in 1934 and the attendant rise in gold price to \$35 per ounce encouraged some rather large-scale attempts to mine the placers in 1935. Heavy equipment (shovels, etc.) was moved in but operations again proved a failure and lasted only briefly (Metzger, 1938, p. 59). All gold operations ceased activity by order of the War

Production Board in 1942 and the district was essentially idle through 1957 (Howard, 1967, p. 123, 124).

The Jicarilla district has been examined and evaluated in considerable detail over the past several decades (Segerstrom and Ryberg, 1974) in an effort to locate precious metal deposits of an economic grade sufficient to sustain mining operations. None of these attempts were successful.

The total mineral production of this district is not known precisely. Gold production prior to 1900 is unverifiable; through 1931 it is inconsequential and is estimated at \$90,000 (Table 1; Lasky and Wooton, 1933). A total of \$72,125 in gold, silver and copper production is credited to the district since then, making a total of \$162,125. Estimated and known base and precious metals production is tabulated in Table

TABLE 1. Precious and base metal production from the Jicarilla mining district, Lincoln County, New Mexico. Compiled from U.S. Bureau of Mines Mineral Yearbooks, Segerstrom and Ryberg (1974) and Johnson (1972). ^aTotal gold placer production is estimated. ^bIncludes some placer silver production.

Year	Ore (short tons)	Copper (lbs)	Gold lode (oz)	Gold placer (oz)	Silver ^b (oz)	Lead (lbs)
Prior to 1902 ^a	---	---	---	4,500	---	---
1902-1911 ^a	---	---	---	646	---	---
1912	---	---	---	48.6	---	---
1914	11,009	387,211	1,861	44	5,376	2,027
1915	---	1,747,090	2,104	63	24,570	---
1916	20,195	1,077,492	1,911	31	7,401	638
1917	15,558	731,894	1,137	218	---	---
1918	6,210	253,405	247	99	---	---
1919	205	4,382	4	---	25	---
1933	130	---	82.62	236.02	50	---
1934	---	---	---	327.81	26	---
1935	---	---	---	309.09	21	---
1936	---	---	---	289.40	20	---
1937	---	---	---	184.2	16	---
1938	---	---	---	162.2	11	---
1939	---	---	---	203	13	---
1940	---	---	---	178	14	---
1941	---	---	---	178	14	---
1942	---	---	---	36	4	---
1946	---	---	---	1.0	---	---
1948	---	---	---	1.0	---	---
TOTAL ^a	53,307	4,201,474	7,347	7,755	37,561	2,665

1. In addition, approximately 8000 tons of iron ore was produced from the district between the periods 1918 to 1921 and 1942 to 1943 (Kelley, 1949; Griswold, 1959).

GEOLOGY

The Jicarilla Mountains consist of an incomplete sequence of Permian through Cretaceous sedimentary rocks (Table 2) that have been intruded by a series of Tertiary igneous intrusives (Fig. 1). Remnants of

TABLE 2. Stratigraphic units in the Jicarilla Mountains (from Budding, 1963, 1964; Segerstrom and Ryberg, 1974). Specific units are described elsewhere in this guidebook.

Recent	Alluvial fan and stream deposits
Pliocene	Ogallala Formation (stream and mudflow deposits)
Oligocene-Eocene	Basaltic dikes Intrusive rocks
Cretaceous	Mesaverde Group Mancos Shale Dakota Group
Triassic	Chinle Formation Santa Rosa Formation
Permian	Artesia Formation(?) San Andres Formation Glorieta Sandstone Yeso Formation

the Ogallala Formation (Tertiary) occur throughout the area (Budding, 1963, 1964). The sedimentary rocks are described elsewhere in this guidebook and the reader is referred to those articles for more information.

The igneous rocks in the Jicarilla Mountains are complex and quite diverse in lithology and composition. More detailed petrographic and geochemical work is needed. Preliminary data (V. T. McLemore, field data, 1990) and reports by Budding (1963, 1964), Ryberg (1968) and Segerstrom and Ryberg (1974) indicate the Jicarilla Mountains contain a lithologic diversity of porphyritic stocks and dikes. Compositions range from syenite to quartz syenite to quartz monzonite to granite (Table 3; De la Roche et al., 1980). The porphyry phenocryst assemblage includes medium-grained plagioclase, hornblende, biotite and quartz. X-ray diffraction studies indicate the groundmass is composed mostly of microcrystalline orthoclase and quartz. The plagioclase is euhedral and present as monomineralic glomerocrysts and single grains. Most phenocrysts have moderate normal zoning. Unzoned phenocrysts are optically sodic-andesine. Alteration in plagioclase is limited to sericite occurring along cleavage planes and at cleavage intersections. Hornblende is euhedral, pleochroic green to brown, and is locally altered to chlorite. Biotite occurs as euhedral to subhedral plates with limited chloritic alteration. Quartz phenocrysts are sparsely distributed as subhedral grains, typically showing resorption surfaces. The groundmass is microgranular orthoclase laths with anhedral, interstitial, quartz grains. In some quartz monzonites and tonolites, xenoliths of foliated, garnet-bearing amphibolite have been observed by Allen and Foord (1991, this volume).

Chemically (Table 3), the porphyries are diverse in being subalkaline to alkaline, and peraluminous but not heavily silica-saturated (8 to 18% normative quartz; Fig. 2). The porphyries plot as trachydacites based on total alkali versus silica (Fig. 3; Le Bas et al., 1986) compared to their extrusive equivalents.

Two age dates of rocks from the Jicarilla Mountains are reported in the literature. Segerstrom and Ryberg (1974) reported a K-Ar age of 37.3 ± 1.5 Ma on biotite from a monzonite. A second age of 38.2 Ma (recalculated using new constants) was reported in Isochron/West (1979, no. 26, p. 25). These ages are among the oldest for Tertiary igneous rocks in Lincoln County, and are similar to some of the volcanic and pluton rocks of the Sierra Blanca complex (Allen and Foord, 1991, this volume).

MINERAL DEPOSITS

Three types of mineral deposits occur in the Jicarilla Mountains: placer, gold-bearing quartz veins and iron skarn/replacement deposits. All three types of deposits have been productive; however, the most significant production has come from the gold placers.

Placer deposits (late Tertiary to Quaternary)

Placer gold occurs in gravels, sands and silts of the Ogallala Formation. Some high-grade gold streaks occur in weathered zones overlying the Tertiary igneous intrusives (Segerstrom and Ryberg, 1974). Gold placers also occur in poorly sorted, unconsolidated fanglomerate and alluvial deposits that are less than a few meters thick. Gold assays are typically low, but range as high as 17 ppm. Gold is less than 2 mm in diameter. The placers consist of gold, scheelite, magnetite, hematite, hornblende, epidote, augite, sphene, apatite, zircon and pyrite (Griswold, 1959; Segerstrom and Ryberg, 1974). The gold is most likely derived from gold-bearing quartz veins as well as gold-rich zones within the intrusives (Johnson, 1972).

Gold-bearing quartz veins

The gold-bearing quartz veins in the Jicarilla district have been classified as Great Plains margin (GPM) deposits by North and McLemore (1986). GPM deposits in New Mexico lie along, or near, the border of the Great Plains with the Southern Rocky Mountains or Basin and Range provinces (McLemore, 1991, this volume) and include gold-bearing breccia pipes and quartz veins, copper and/or lead/zinc skarns and iron skarns. They are typically associated with alkalic rocks and with de-

TABLE 3. Chemical analyses of selected samples from Jicarilla Mountains. Samples J22.1, J27.2, J27.3 collected by senior author and analyzed by Chris McGeer (NMBM&MR, X-ray Laboratory). Samples J1, J11, J15, J17, J19, J53 from Segerstrom and Ryberg (1974). ^aTotal iron reported as Fe₂O₃. ^bLithology from discriminant diagram of De la Roche et al. (1980).

Wt %	J22.1	J27.2	J27.3	J27.4	J1	J11	J15	J17	J19	J53
SiO ₂	65.4	54.5	63.5	70.4	64.2	64.6	56.0	57.7	61.8	60.9
TiO ₂	0.51	0.54	0.53	0.52	0.47	0.26	0.77	0.72	0.82	0.70
Al ₂ O ₃	16.4	16.5	16.0	16.8	15.8	17.5	15.9	15.8	16.7	16.2
Fe ₂ O ₃	3.34 ^a	3.36 ^a	3.9 ^a	1.78 ^a	2.3	2.2	4.7	4.8	2.0	3.4
FeO	--	--	--	--	2.3	0.84	2.8	3.1	2.6	2.8
MnO	0.06	0.05	0.06	0.01	0.06	--	0.10	0.12	0.07	0.06
MgO	1.09	1.09	1.21	0.47	1.7	0.57	2.00	3.7	1.8	2.6
CaO	4.5	3.41	3.67	0.63	2.20	0.89	3.7	4.0	3.3	3.9
Na ₂ O	4.6	5.97	6.07	6.49	4.80	4.90	5.70	5.3	5.4	4.7
K ₂ O	2.86	3.18	3.25	1.14	3.00	6.80	3.00	2.9	3.2	3.0
P ₂ O ₅	0.32	0.32	0.32	0.31	0.27	0.14	0.53	0.5	0.37	0.23
LOI	0.95	0.78	1.51	2.12	--	--	--	--	--	--
H ₂ O ⁺	--	--	--	--	1.5	0.9	1.2	0.82	0.45	1.1
H ₂ O ⁻	--	--	--	--	0.59	0.40	0.15	0.38	0.55	0.42
CO ₂	--	--	--	--	0.58	0.08	3.4	0.05	0.34	0.05
TOTAL	100.03	99.99	100.02	100.67	99.77	100.08	99.95	99.89	99.40	100.06
CIPW NORM										
Apatite	0.71	0.71	0.70	0.66	0.60	0.31	1.71	1.30	0.83	0.52
Ilmenite	0.98	1.03	1.01	0.95	0.91	0.49	1.47	1.40	1.60	1.36
Magnetite	2.44	2.66	2.83	1.24	3.38	1.95	6.86	7.14	2.98	5.05
Orthoclase	16.99	18.84	19.16	6.4	17.94	40.05	17.81	17.54	19.35	18.13
Albite	39.14	50.67	51.24	52.75	41.11	41.34	48.46	45.92	46.78	40.67
Anorthoclase	15.74	8.86	7.80	1.06	9.26	3.49	8.98	11.01	12.16	14.57
Diopside	3.97	5.17	6.90	--	5.17	7.98	5.32	5.28	1.84	3.23
Hyperssthene	1.27	1.00	0.54	0.83	3.46	0.93	1.95	5.11	3.28	4.01
Corundum	--	--	--	8.63	--	6.58	--	2.64	--	1.59
Quartz	17.85	10.27	8.26	25.97	7.79	7.70	3.18	4.11	9.24	10.94
Hornblende	0.30	0.85	--	--	--	--	--	--	--	--
TOTAL	99.08	99.21	98.51	98.85	97.10	98.70	95.20	98.64	98.06	98.49

posits of iron, molybdenum, rare-earth elements, fluorite, tungsten and niobium.

Quartz veins in the Jicarilla district strike predominantly north-westward, but some strike north-south or northeastward and all are steeply dipping. Minerals associated with the veins include hematite, pyrite (rarely arsenopyrite), chalcopyrite, sphalerite, galena and gold (Lindgren et al., 1910; Segerstrom and Ryberg, 1974).

Additionally, some areas contain disseminations of gold and quartz stockworks within the igneous intrusives. Silicification is common and sericitization occurs locally. Veins range in width up to 10 m (Lindgren et al., 1910), but very little information derived during the lode mining period (ca. 1880–1910) has been preserved and little is known regarding the extent and grade of these deposits.

Iron-skarn/replacement and vein deposits

Numerous iron-skarn/replacement and vein deposits occur throughout Lincoln County and are spatially associated with Tertiary igneous intrusives (Kelley, 1949; Harrer and Kelly, 1963; Griswold, 1959; McLemore, 1991, fig. 1). The deposits in the Jicarilla Mountains are

small and low grade (less than 40% Fe). They consist of a series of iron-ore replacement bodies and veins in limestones of the San Andres or Yeso Formations. Ore minerals consist of magnetite and hematite in a gangue of calcite, quartz, fluorite, and locally pyrrhotite, chalcopyrite and purite (T. B. Thompson, written comm., 1991). Pyroxene, tremolite, actinolite, epidote and phlogopite are also present. The presence of calc-silicate minerals and proximity of the iron-ore deposits to the igneous intrusives suggest a magmatic origin for the mineralization. However, detailed petrographic, mineralogic and geochemical data are needed to determine the origin of these deposits.

DISCUSSION

The Tertiary intrusives of the Jicarilla Mountains range in age up to at least 37–38 Ma (younger intrusives are present but have not been dated). These rocks represent part of the early stage of Tertiary magmatic activity that ultimately formed the Lincoln County porphyry belt. Some of the Jicarilla intrusives are subalkalic to alkalic (Fig. 3); however, not all of the intrusives in the Jicarilla Mountains have been studied. A younger magmatic event occurred in Lincoln County around 28–26




FIGURE 2. Variation diagrams of oxides versus silica for samples from Jicarilla district. Chemical analyses are in Table 3.

Ma and is represented by the Capitan pluton and younger rocks of the Sierra Blanca igneous complex (Thompson, 1972; Allen and Foord, 1991, this volume). These two magmatic events may represent two separate events or they may represent one single period of magmatic activity. There are not enough geochronologic data to distinguish between a single long event or two separate events in the Lincoln County porphyry belt. A detailed geochronologic study in this area would provide the information needed to answer this question.

The mineral deposits in the Jicarilla Mountains are similar to other mineral deposits in the western United States that occur near or along the margin of the tectonically stable Great Plains and the more tectonically active Basin-and-Range and Rocky Mountains provinces. Tertiary deposits in New Mexico that occur along this margin have been called Great Plains margin (GPM) deposits (North and McLemore, 1986; McLemore, 1991, this volume). The intersection with the west-northwest-trending Capitan lineament appears to have periodically leaked magma and perhaps controlled mineralization in Lincoln County since about 33 Ma (McLemore, 1991).

A magmatic origin for these deposits is unproven, but there is evi-




FIGURE 3. Plot of Na₂O + K₂O versus silica for samples from Jicarilla district. Chemical analyses are in Table 3.

dence to support such an origin. The close proximity of iron-skarn/replacement deposits and gold-bearing quartz veins to the Jicarilla intrusives suggests a magmatic derivation. The presence of some gold placers in the weathered zones on top of the Jicarilla intrusives suggests that gold may be disseminated within the intrusives. Other evidence from other GPM deposits supporting a magmatic origin is summarized by Allen and Foord (1991, this volume) and McLemore (1991, this volume).

The origin of the mineral deposits in the Jicarilla district and elsewhere in Lincoln County is complex, as it is elsewhere along the Great Plains margin. The diversity in lithology, composition and age of the intrusives indicates a complex process of differentiation and fractionation of magmas, perhaps involving upper mantle and lower crustal sources. Much work remains to be done, including geochemical, isotopic and geochronological studies.

MINERAL RESOURCE POTENTIAL

The Jicarilla district is known to host a placer gold resource, perhaps the largest remaining in New Mexico (about 5.4 million yd³), but the grade is low (0.02–0.04 oz/yd; Segerstrom and Ryberg, 1974). However, due to the low grade and scarcity of water, this resource is uneconomic in today's market. The widely disseminated placer-gold deposits might be indicative of a buried mineralized deposit. This potential has not been investigated by drilling and is speculative. GPM deposits locally host gold-bearing breccia deposits, such as at Ortiz (Maynard et al., 1990). Similar deposits could exist in the Jicarilla district, but detailed mapping followed by subsurface drilling are required to verify this.

The iron deposits are small and low grade and are not economic in today's market. Certainly if a small steel mill were built in Lincoln County, these deposits should be re-examined for their iron resource.

Occasionally, the Jicarilla district has been promoted more upon the supposed presence of platinum group metals (PGMs) than upon gold. The NMBM&MR has analyzed many samples from this district over the last 30 yrs or so. No PGMs have ever been detected in samples from the district (McLemore et al., 1989). Economic deposits of PGMs typically occur in ultramafic rocks; these rocks have not been identified from the Jicarilla district. Although alkalic rocks similar to those found in the Jicarilla district may have the potential for the occurrence of PGMs, the potential for economic concentrations from this area is poor (McLemore et al., 1989).

ACKNOWLEDGMENTS

Many thanks to Tommy B. Thompson and George Ryberg for reviewing an earlier version of this manuscript. Darren Dresser is acknowledged for assistance in sample preparation and in compiling Fig. 1.

REFERENCES

- Allen, M. S. and Foord, E. E., 1991, Geochemical and isotopic characteristics of the Lincoln County porphyry belt, New Mexico: implications for regional tectonics and mineral deposits: *New Mexico Geological Society, Guidebook* 42.
- Anderson, E. C., 1957, The metal resources of New Mexico and their economic features through 1954: *New Mexico Bureau of Mines and Mineral Resources, Bulletin* 39, 183 p.
- Budding, A. J., 1963, Origin and age of superficial structures, Jicarilla Mountains, central New Mexico: *Geological Society of America Bulletin*, v. 74, p. 203–208.
- Budding, A. J., 1964, Geologic outline of the Jicarilla Mountains, Lincoln County, New Mexico: *New Mexico Geological Society, Guidebook* 15, p. 82–86.
- De la Roche, H., Leterrier, J., Grandclaude, P. and Marchal, M., 1980, A classification of volcanic and plutonic rocks using R_1R_2 -diagram and major-element analyses—its relationship with current nomenclature: *Chemical Geology*, v. 29, p. 183–210.
- Griswold, G. B., 1959, Mineral deposits of Lincoln County, New Mexico: *New Mexico Bureau of Mines and Mineral Resources, Bulletin* 67, 117 p.
- Harrer, C. M. and Kelly, F. J., 1963, Reconnaissance of iron resources in New Mexico: U.S. Bureau of Mines, *Information Circular* 8190, 112 p.
- Henderson, C. W., 1908, Gold, silver, copper, lead, and zinc in the western states—New Mexico: *Mineral Resources of the United States*, U.S. Geological Survey, p. 506–519.
- Henderson, C. W., 1910, Gold, silver, copper, lead, and zinc in the western states—New Mexico: *Mineral Resources of the United States*, U.S. Geological Survey, p. 534–552.
- Howard, E. V., 1967, Metalliferous occurrences in New Mexico: *State Planning Office*, Santa Fe, 270 p.
- Irvine, T. N. and Baragar, W. R. A., 1971, A guide to the chemical classification of the common volcanic rocks: *Canadian Journal of Earth Sciences*, v. 8, p. 523–548.
- Johnson, M. G., 1972, Placer gold deposits of New Mexico: U.S. Geological Survey, *Bulletin* 1348, 46 p.
- Jones, F. A., 1904, *New Mexico Mines and Minerals: World's Fair Edition*, Santa Fe, 349 p.
- Kelley, V. C., 1949, *Geology and economics of New Mexico iron-ore deposits*: University of New Mexico, *Publications in Geology*, no. 2, 246 p.
- Lasky, S. G. and Wootton, T. P., 1933, The metal resources of New Mexico and their economic features: *New Mexico Bureau of Mines and Mineral Resources, Bulletin* 7, 178 p.
- Le Bas, M. J., Le Maitre, R. W., Streckeisen, A. and Zanettin, B., 1986, A chemical classification of volcanic rocks based on the total alkali-silica diagram: *Journal of Petrology*, v. 27, p. 745–750.
- Lindgren, W., Graton, L. C. and Gordon, C. H., 1910, The geology and ore deposits of New Mexico: U.S. Geological Survey, *Professional Paper* 68, 361 p.
- Maynard, S. R., Nelson, C. J., Martin, K. W. and Schultz, J. L., 1990, Geology and gold mineralization of the Ortiz Mountains, Santa Fe County, New Mexico: *Mining Engineering*, August, p. 1007–1011.
- McLemore, V. T., 1991, Base- and precious-metals deposits in Lincoln and Otero Counties, New Mexico: *New Mexico Geological Society, Guidebook* 42.
- McLemore, V. T., Eveleth, R. W., Brandvold, L. A. and Robertson, J. M., 1989, Platinum-group metals in New Mexico: *New Mexico Geology*, p. 29–30, 33.
- Metzger, O. H., 1938, Gold mining in New Mexico: U.S. Bureau of Mines, *Information Circular* 6987, 71 p.
- North, R. M. and McLemore, V. T., 1986, Silver and gold occurrences in New Mexico: *New Mexico Bureau of Mines and Mineral Resources, Resource Map* 15, 32 p.
- Ryberg, G. E., 1968, The geology of the Jicarilla Mountains, Lincoln County, New Mexico [M.S. thesis]: Albuquerque, University of New Mexico, 95 p.
- Segerstrom, K. and Ryberg, G. E., 1974, Geology and placer-gold deposits of the Jicarilla Mountains, Lincoln County, New Mexico: U.S. Geological Survey, *Bulletin* 1308, 25 p.
- Thompson, T. B., 1972, Sierra Blanca igneous complex, New Mexico: *Geological Society of America Bulletin*, v. 83, p. 2341–2356.
- Wright, P. E., 1932, The Jicarilla mining district of New Mexico: *Mining Journal*, v. 16, no. 8, p. 7.