



Paleomagnetic studies in the Jemez Mountains, New Mexico: A progress report on Quaternary volcanic rocks from Valles Caldera VC-2A, Sulphur Springs, and Lower Permian strata in San Diego Canyon and from VC-2B

John W. Geissman and Hope J. Mullally, 1996, pp. 115-120

in:

Jemez Mountains Region, Goff, F.; Kues, B. S.; Rogers, M. A.; McFadden, L. S.; Gardner, J. N.; [eds.], New Mexico Geological Society 47th Annual Fall Field Conference Guidebook, 484 p.

This is one of many related papers that were included in the 1996 NMGS Fall Field Conference Guidebook.

Annual NMGS Fall Field Conference Guidebooks

Every fall since 1950, the New Mexico Geological Society (NMGS) has held an annual [Fall Field Conference](#) that explores some region of New Mexico (or surrounding states). Always well attended, these conferences provide a guidebook to participants. Besides detailed road logs, the guidebooks contain many well written, edited, and peer-reviewed geoscience papers. These books have set the national standard for geologic guidebooks and are an essential geologic reference for anyone working in or around New Mexico.

Free Downloads

NMGS has decided to make peer-reviewed papers from our Fall Field Conference guidebooks available for free download. Non-members will have access to guidebook papers two years after publication. Members have access to all papers. This is in keeping with our mission of promoting interest, research, and cooperation regarding geology in New Mexico. However, guidebook sales represent a significant proportion of our operating budget. Therefore, only *research papers* are available for download. *Road logs, mini-papers, maps, stratigraphic charts*, and other selected content are available only in the printed guidebooks.

Copyright Information

Publications of the New Mexico Geological Society, printed and electronic, are protected by the copyright laws of the United States. No material from the NMGS website, or printed and electronic publications, may be reprinted or redistributed without NMGS permission. Contact us for permission to reprint portions of any of our publications.

One printed copy of any materials from the NMGS website or our print and electronic publications may be made for individual use without our permission. Teachers and students may make unlimited copies for educational use. Any other use of these materials requires explicit permission.

This page is intentionally left blank to maintain order of facing pages.

PALEOMAGNETIC STUDIES IN THE JEMEZ MOUNTAINS, NEW MEXICO: A PROGRESS REPORT ON QUATERNARY VOLCANIC ROCKS FROM VALLES CALDERA VC-2A, SULFUR SPRINGS, AND LOWER PERMIAN STRATA IN SAN DIEGO CANYON AND FROM VC-2B

JOHN WM. GEISSMAN¹ and HOPE J. MULLALLY²

¹Department of Earth and Planetary Sciences, University of New Mexico, Albuquerque, NM 87131-1116;

²804 Ortiz Drive SE, Apt. 2, Albuquerque, NM 87108

Abstract—Paleomagnetic data from Quaternary volcanic rocks intersected in the VC-2A drilling experiment are interpreted to indicate that, despite considerable alteration in an elevated geothermal gradient, the initial, reverse polarity thermoremanent magnetization of intracaldera and pre-caldera ignimbrites is preserved over much of the 1731 ft sampled interval. In all volcanic materials, magnetite is the principal remanence carrier. Overall, the data are consistent with the interpreted stratigraphy, but many rocks below about 800 ft are partially to completely overprinted, due to hydrothermal alteration, by a positive inclination (normal polarity) remanence interpreted to be less than 780 ka (Brunhes normal polarity chron). Below about 1400 ft, NRM intensities decrease by about an order of magnitude, to between 2 and 7 mA/m. Lower Permian Abo and Yeso Formation strata from the VC-2B experiment yield steep, positive inclination magnetizations as the single or dominant component, suggesting that these rocks have been remagnetized during the Brunhes. The mechanism of remagnetization may involve a combination of heating to no greater than 300°C and changes to the population of hematite grains. The same Permian strata exposed in San Diego Canyon, however, do not show evidence of partial remagnetization related to Quaternary magmatism. Of 16 sites fully demagnetized, 12 yield high-precision results providing a grand mean, corrected for gentle tilt of the strata (Declination= 148°, Inclination= -1.4°, $\alpha_{95} = 4.6^\circ$, $k = 88.7$; paleomagnetic pole at 44N, 120E, $dp = 2.3^\circ$, $dm = 4.6^\circ$), that is displaced in a small, and statistically indistinguishable clockwise sense ($R = 2.6^\circ$, $\Delta R = 3.7^\circ$) from an expected Early Permian direction. The San Diego Canyon data are interpreted to indicate that this part of the westernmost margin of the Rio Grande rift did not experience appreciable cumulative vertical axis rotation since the Early Permian.

INTRODUCTION

The Jemez Mountains and its Miocene to Quaternary volcanism have been the focus of considerable paleomagnetic research. In the early 1960s, fundamental paleomagnetic and geochronologic information was obtained from Valles caldera rocks (Doell et al., 1968) that served as the basis for the initial geomagnetic polarity time scale and defined some of the fine structure of that scale (e.g., identification of the Jaramillo "event" or sub-chron by Doell and Dalrymple, 1966). Three drilling experiments in the Valles Caldera, as part of the Continental Scientific Drilling Program, have provided new opportunities for investigation.

Experiment VC-1 (Fig. 1) was drilled in the southwest moat of the Valles caldera and penetrated post-Bandelier volcanic rocks and upper Paleozoic strata. Paleomagnetic and rock magnetic work on VC-1 (Geissman, 1988) demonstrated that upper Paleozoic strata had been uniformly partially remagnetized during Quaternary volcanism. The thermal demagnetization characteristics of these secondary magnetizations were used to better understand the thermal potency of viscous partial thermoremanent magnetization acquisition. Paleomagnetic data from the Quaternary volcanic part of the core stratigraphy aided in the identification of the volcanic materials intersected. Experiment VC-2A (Figs. 1, 2) was drilled into the Sulfur Springs hydrothermal system, near the intersection of the ring fracture zone and the western margin of the resurgent dome and principally intersected intercaldera and pre-caldera ignimbrites. VC-2B, cored 0.5 km northeast of VC-2A (Fig. 1) was located with the intent of penetrating well into the floor of the caldera, and intersected Quaternary volcanic rocks and over 3000 ft of upper Paleozoic strata and Proterozoic rocks.

This contribution is a progress report on Quaternary volcanic rocks from the VC-2A and Permian strata from the VC-2B experiments. Study of VC-2A aids in defining the volcanic stratigraphy and in evaluating the degree to which the rocks may have been remagnetized during hydrothermal alteration at moderate temperatures. Interest in the Permian section of VC-2B centers around the ability of these rocks to retain a "primary" Early Permian magnetization. Here we also present the results of preliminary work on a well-exposed section of Permian strata in San Diego Canyon which became important because initial data from Permian strata from VC-2B indicated that these rocks were pervasively remagnetized, prob-

ably in the late Quaternary. To test the hypothesis that remagnetization was in direct response to Quaternary magmatism, information on the intensity and laboratory unblocking temperature spectra of Permian strata, as unaltered and thermally unaffected as possible, was required. Research on all three components of this study is still in progress.

GENERAL GEOLOGY AND SAMPLING

Volcanic and volcaniclastic rocks, and alteration features, intersected in the VC-2A experiment (total depth of 1731 ft) at Sulfur Springs (Figs.




FIGURE 1. Location map of the Valles caldera and VC-2A, VC-1 and VC-2B core holes. Stippled pattern shows area of intense near-surface leaching. Modified from Goff et al. (1987).




FIGURE 2. Generalized stratigraphy of VC-2A, Valles caldera, New Mexico, modified from Goff et al. (1987), showing locations of core segments (samples) collected.

1, 2) have been described in detail (e.g., Goff et al., 1987; Starquist, 1988; Goff and Gardner, 1994). The section is comprised principally of the intracaldera Tshirege and Otowi Members and, possibly pre-caldera ignimbrites. Intracaldera ignimbrites would have formed at about 1.13 and 1.50 Ma, during the Matuyama reversed polarity chron. The age of pre-caldera ignimbrites intersected at about 1550 ft in VC-2A is unknown, but they may correlate with San Diego Canyon ignimbrites dated at about 1.78 Ma (Hulen et al., 1992). The VC-2A experiment revealed that the active hydrothermal system at Sulfur Springs consists of a thin (about 6 ft) acid condensation zone above a vapor-dominated zone (much of the section) and a liquid-dominated zone at the bottom. The unequilibrated bottom hole temperature was 212°C. All of the rocks in VC-2A have been moderately to intensely altered. The presence of illite throughout the section was interpreted by Goff et al. (1987) and WoldeGabriel and Goff (1989) to indicate that the modern acid sulfate environment is superimposed on an older, higher temperature alteration. Below about 450 ft, most materials contain abundant illite, chlorite, calcite, quartz, pyrite and fluorite. A total of 115 azimuthally unoriented sections (Fig. 2) of core from VC-2A were obtained from Los Alamos National Laboratory in 1987. Because core material between about 360 and 780 ft is uniform in character, we deferred sampling this interval for our initial efforts. The core sections, here referred to as independent samples, range from about 2 in. to over 2 ft in length. The inclination of the VC-2A borehole is within a few degrees of vertical throughout the experiment (Goff et al., 1987). The hole was spudded vertically and the requirement of extremely high core recovery mandated that drilling was carried out slowly to maintain a constant borehole orientation.

The VC-2B (Fig. 1) experiment was continuously cored to a total depth

of 5780 ft and penetrated about 2500 ft of Permian and Pennsylvanian carbonate and detrital strata between a depth of about 2600 ft and 5100 ft (Hulen and Gardner, 1989). This section is overlain by intracaldera and pre-caldera volcanic rocks and volcaniclastic strata. Of the more than 300 azimuthally unoriented sections of core collected from VC-2B for paleomagnetic study, 79 are from Lower Permian Abo and Yeso strata between 2620 ft and 4250 ft. The core sections (independent samples) range from about 2 in. to over 2 ft in length. Orientation surveys through about 4600 ft of the section indicate that the VC-2B borehole is less than one degree from vertical (Hulen and Gardner, personal commun., 1989). Cross-beds and laminar beds have dips up to about 30° (Hulen and Gardner, 1989). Yeso and Abo rocks are largely unaltered, except adjacent to structures or in high permeability, coarse-grained sandstones.

Lower Permian strata, principally of the Abo and Yeso Formations, are exceptionally well-exposed in San Diego Canyon (Smith et al., 1970). Romer (1960) reported the age of the Abo Formation as Wolfcampian based on vertebrate paleontology, and Needham and Bates (1943) placed the overlying Yeso strata into the lower two thirds of the Leonardian Series. We sampled 53 discrete beds (paleomagnetic sampling sites, with site number increasing stratigraphically upward) and the lowermost, moderately welded part of the Otowi Member of the Bandelier Tuff along a section about 2.5 mi south of Jemez Springs below Virgin Mesa. At each bed, between 7 and 15 independent samples, as drilled cores, were obtained and, where possible, the orientation of bed contacts was also measured.

LABORATORY METHODS

Procedures used for the preparation of individual specimens from drill core segments for remanence and rock magnetic measurements, progressive demagnetization, and statistical evaluation of demagnetization data follow standard practices, as detailed in Geissman (1988). The nature of the natural remanent magnetization (NRM) in the rocks discussed was evaluated using progressive alternating field or thermal demagnetization, and in some cases a combination of both methods. Directions of individual magnetization "components" contributing to the NRM were determined following the method of Kirschvink (1980); unanchored line segments, defined by at least four demagnetization steps, with MAD values less than 10° were accepted for further analysis. For azimuthally unoriented core, only the inclination of a magnetization component, generally relative to the present horizontal, can be confidently determined. In examples of demagnetization results (Figs. 3, 6, 7), the demagnetization results are plotted according to the method of Roy and Park (1974), where the vertical component of the magnetization is plotted against the (full) horizontal component, facilitating measurement of the inclination of the magnetization isolated over a range of demagnetization steps.

PALEOMAGNETIC DATA

Quaternary rocks, VC-2A

Most Quaternary volcanic and volcaniclastic rocks intersected in VC-2A yield interpretable results in either progressive alternating field or thermal demagnetization (Fig. 3). One common demagnetization behavior involves the removal of essentially a single component of magnetization over a wide range of high peak alternating fields and laboratory unblocking temperatures exceeding 350°C (e.g., core sample 223-6, Fig. 3). In these cases, inclinations of the magnetization isolated are usually of moderate negative values. Another behavior encountered involves the removal of a substantial steep to moderate positive inclination magnetization at low peak fields or laboratory unblocking temperatures and the isolation of, again, a moderate negative inclination magnetization above peak fields of about 30 mT and laboratory unblocking temperatures of about 350 to 400°C (core samples 52-5 and 45-3, Fig. 3). Finally, many core samples from depths greater than about 800 ft exhibit more complex behavior, such as (1) alternating field and thermal demagnetization do not yield the same results (core sample 355-4, Fig. 3); (2) the magnetization isolated is of unexpectedly shallow inclination; or (3) a magnetization of positive inclination is the principal magnetization and is isolated over relatively low alternating fields and/or laboratory unblocking temperatures. In all thermal demagnetization experiments, over 99% of




FIGURE 3. Representative modified orthogonal demagnetization diagrams (Zijderveld, 1967; Roy and Park, 1974) for specimens from samples from VC-2A, arranged in stratigraphic order. In each diagram, the endpoint of the magnetization vector measured after each step of demagnetization is simultaneously projected onto the (artificial with unoriented core) horizontal plane (solid symbols) and the true vertical plane (open symbols) assuming the axis of the core segment is vertical. Peak demagnetization inductions or temperatures are given along the vertical projections. Magnetization intensity (in milliamperes/meter, mA/m) is indicated by a tic along one axis. Modified orthogonal demagnetization diagrams give the true inclination of a single magnetization vector isolated over a series of several demagnetization steps.

the magnetization is unblocked by about 580°C. NRM intensities of the core samples over most of the sampled section range from 3 to 10 mA/m (Fig. 4). At depths greater than about 1450 ft, NRM intensities decrease by about an order of magnitude.

The distribution of values of inclination of the magnetization isolated over the highest peak fields and laboratory unblocking temperatures (Fig. 5) is complicated. With the exception of the section above 360 ft, where the inclinations are consistently negative (presumably reversed polarity magnetizations), there no strong correlation with rock type.

Lower Permian strata, VC-2B

Paleomagnetic work on samples from Lower Permian strata in VC-2B is in preliminary stages. All rocks analyzed to date have a moderate positive inclination NRM, and the principal magnetization, isolated beginning at less than 100°C up to maximum unblocking temperatures of hematite (about 680°C), is of steep to moderate positive inclination. Assuming this magnetization is north-seeking, it can be interpreted as of normal polarity (Fig. 6). Less than 20% of the 35 core samples fully demagnetized to date contain a sub-horizontal inclination magnetization (e.g., sample 453B, Fig. 6) isolated at temperatures above 600°C or exhibit a progressive shallowing of inclination at unblocking temperatures above about 500°C. The average NRM intensity for 66 specimens from 52 core segments is 25.8 mA/m (std. dev. = 17.9 mA/m).




FIGURE 4. NRM intensity vs. core sample depth for Quaternary rocks from the VC-2A experiment.




FIGURE 5. Inclination of the magnetization isolated, toward the origin, over the highest peak fields or laboratory unblocking temperatures vs. core sample depth for Quaternary rocks from the VC-2A experiment.




FIGURE 6. Representative modified orthogonal demagnetization diagrams (Zijderveld, 1967; Roy and Park, 1974) for specimens from Permian strata intersected in VC-2B. Conventions follow those in Figure 3.




FIGURE 8. Partial equal area projection of directions of site means (small circles) in geographic coordinates, from 16 of the 53 sites sampled (the other 37 have not been demagnetized) in the Permian section in San Diego Canyon. Four of the site means (circles cut by dashed lines) have been rejected because of unacceptably high within-site dispersion. The grand mean of the 12 accepted means is given by the large circle (full for geographic coordinates and broken for stratigraphic coordinates). "Expected", time-averaged Early to Late Permian (earliest Triassic) field directions, all of shallow negative inclination, are shown as squares. Expected directions are derived from paleomagnetic poles in Van der Voo (1989) (305 Ma), Irving and Irving (1982) (300 Ma), and Miller and Kent (1988) (301, 281, 261 and 248 Ma).

activation theory for single domain grains (e.g., Pullaiah et al., 1975; Walton, 1980) full thermal remagnetization of hematite over the last 780 ka of normal field polarity requires temperatures exceeding about 600°C, and there is no evidence that the Permian strata were subjected to such high temperatures, at even a localized (e.g., vein or brecciated zone) scale. Homogenization temperatures for primary fluid inclusions in minerals from deep veins cutting Permian and older strata are similar to equilibration temperatures measured following the experiment (Hulen and Gardner, 1989), and the highest fluid inclusion temperatures measured in VC-2B material are about 310°C (Goff and Gardner, 1994). Ito and Tanaka (1995) reported zircon fission-track dates for Permian strata between 450 and 600 Ma, which they interpreted to indicate that the detrital zircons originated from Precambrian rocks and were partially annealed. These dates are inconsistent with pervasive thermal annealing at high temperatures at any time after the Permian. Finally, clay fractions extracted from the Permian strata contain mixed-layer illite/smectite, with up to 45% expandable interlayers (Hulen and Gardner, 1989). Such a high concentration of expandable interlayers, Hulen and Gardner (1989) argued, would not survive present thermal conditions and must reflect the relatively impermeable nature of the Permian beds and their isolation from K-bearing fluids. If current thermoviscous activation theories for hematite-dominated assemblages are realistic approximations to natural conditions, then a tentative explanation for remagnetization must involve some form of reconstitution of extant hematite and moderate temperatures or the growth of abundant authigenic hematite in these rocks over the past 780 ka.

For Lower Permian strata in San Diego Canyon, assessing the antiquity of a well-grouped and well-defined magnetization in essentially flat-lying redbeds is not straightforward. The uniform SSE declination and generally shallow inclination (reverse polarity) of the magnetization in all interpretable demagnetization results is consistent with a primary, Early Permian age of magnetization acquisition. The preliminary grand mean for 12 sites (Declination = 148°, Inclination = -1°) is similar to the result

(Declination = 152°, Inclination = -6°, $N = 74$ samples, $\alpha_{95} = 1.6^\circ$) reported by Steiner (1988) for the Abo Formation exposed at Abo Pass, along the eastern margin of the Rio Grand Rift. It is difficult to evaluate the significance of the difference in directions because the mean direction determined by Steiner (1988) was based on independent sample directions, not "site" (bed) mean directions. For the San Diego Canyon strata, none of the beds, including those within 20 m of the unconformity with the overlying Bandelier Tuff, show evidence of partial remagnetization during emplacement of the reverse polarity Bandelier ignimbrites. They do not contain a well-defined reverse polarity magnetization unblocked up to moderate (e.g., 300 to 500°C) temperatures, as was seen in Permian rocks from the VC-1 experiment (Geissman, 1988). Assuming that the magnetization characteristic of the Lower Permian strata is primary, comparison with expected Early Permian field directions (Fig. 8) reveals that the San Diego Canyon result is statistically indistinguishable from expected directions. This preliminary compilation may imply a small ($R = 2.6^\circ$, $\Delta R = 3.7^\circ$; using the estimated 281 Ma paleomagnetic pole of Miller and Kent, 1988), yet statistically indistinguishable clockwise, cumulative vertical axis rotation of this part of the easternmost margin of the Colorado Plateau since the Permian.

CONCLUSIONS AND FUTURE WORK

Preliminary paleomagnetic data from the VC-2A and VC-2B experiments and a well-exposed section of Permian strata in San Diego Canyon contribute to an improved understanding of the thermochemical history of rocks within the eastern margin of the Valles caldera. The retention of a reverse polarity magnetization in the Quaternary volcanic section in VC-2A supports the interpretation that the rocks intersected are largely intracaldera deposits. Many materials appear to have been relatively immune to intense hydrothermal alteration and despite the complexities of the data from VC-2A, we conclude that the primary thermoremanent magnetizations carried in the appropriate assemblages of magnetite grains can persist in extreme vapor- and hot water-dominated hydrothermal systems. Until more data are obtained from critical depth intervals, we cannot evaluate the timing of remanence acquisition in parts of the section relative to possible localized tilting or slumping.

Permian strata in San Diego Canyon yield, for the most part, high-quality paleomagnetic data, consistent with an Early Permian age of acquisition. Future work on these rocks will involve completion of the additional 37 sites collected and attempts to involve field-based tests, at least evaluating along-strike integrity of the magnetization of selected beds, to better define the age of the well-grouped magnetization in these rocks. Similar Permian strata intersected in VC-2B are partially to seemingly completely remagnetized and exactly how remagnetization occurred is not clear. One possibility is that hydrothermal alteration led to changes in the hematite mineralogy in these rocks and modification of the blocking temperature spectra for these materials. Thermoviscous processes, as conventionally understood for single-domain hematite, are insufficient to result in complete remagnetization. Further paleomagnetic work will focus on determining the level of remagnetization through the Permian section and inspecting, in greater detail, the unblocking characteristics of these rocks between about 630 and 690°C.

ACKNOWLEDGMENTS

Thanks to Jamie Garner and Fraser Goff for incessant encouragement during this study and access to VC-2A and VC-2B core, and to Roberto Molina-Garza for assistance in the laboratory. Steve Harlan and David Iseri aided in the sampling of VC-2A and VC-2B core, and Gary Acton, Dana Bahar, Lucinda Geissman, Roberto Molina-Garza, Sean Mullally, Katerina Petronotis and Cathy Ratcliff assisted in sampling surface exposures in San Diego Canyon. The first author is responsible for all of the paleomagnetic work on VC-2A material. The second author performed all measurements on the San Diego Canyon rocks. Both authors contributed to work on the VC-2B core. Aspects of Hope Mullally's research were supported by the Department of Earth and Planetary Sciences, UNM, and a Geological Society of America Graduate Research Award. Funding for generic paleomagnetic investigations in New Mexico from the New Mexico Bureau of Mines and Mineral Resources is greatly

appreciated. At short notice, Bill McIntosh and Fraser Goff provided excellent reviews of the manuscript and the authors appreciate their suggestions for improving the contribution. We thank Barry Kues for catching an important error in an early draft of the manuscript.

REFERENCES

- Butler, R. F., 1992, Paleomagnetism: magnetic domains to geologic terranes: Oxford, England, Blackwell Scientific, 319 p.
- Doell, R. R. and Dalrymple, G. B., 1966, Geomagnetic polarity epochs—a new polarity event and the age of the Brunhes Matuyama boundary: *Science*, v. 152, p. 1060–1061.
- Doell, R. R., Dalrymple, G. B., Smith, R. L. and Bailey, R. A., 1968, Paleomagnetism, potassium-argon ages, and geology of rhyolites and associated rocks of the Valles caldera, New Mexico: Geological Society of America, *Memoir* 116, p. 211–248.
- Geissman, J. W., 1988, Paleomagnetism and rock magnetism of Quaternary volcanic rocks and late Paleozoic strata, VC-1 core hole, Valles Caldera, New Mexico, with emphasis on remagnetization of late Paleozoic strata: *Journal of Geophysical Research*, v. 93, p. 6001–6025.
- Goff, F. and Gardner, J. N., 1994, Evolution of a mineralized geothermal system, Valles caldera, New Mexico: *Economic Geology*, v. 89, p. 1803–1832.
- Goff, F., Neilson, D. L., Gardner, J. N., Hulen, J. B., Lysne, P., Shevenell, L. and Rowley, J. C., 1987, Scientific drilling at Sulfur Springs, Valles caldera, New Mexico: core hole VC-2A: *Eos, Transactions of the American Geophysical Union*, v. 64, p. 661–662.
- Hulen, J. B. and Gardner, J. N., 1989, Field geologic log for Continental Scientific Drilling Program corehole VC-2B, Valles caldera, New Mexico: Earth Science Laboratory, University of Utah Research Institute, Salt Lake City, 92 p.
- Hulen, J. B., Nielson, D. L. and Little, T. M., 1991, Evolution of the western Valles caldera complex, New Mexico: evidence from intracaldera sandstones, breccias, and surge deposits: *Journal of Geophysical Research*, v. 96, p. 8127–8142.
- Irving, E. and Irving, G. A., 1982, Apparent polar wander paths from Carboniferous through Cenozoic and the assembly of Gondwana: *Geophysical Surveys*, v. 5, p. 141–188.
- Kirschvink, J. L., 1980, The least-squares line and plane and the analysis of paleomagnetic data: *Geophysical Journal of the Royal Astronomical Society*, v. 62, p. 699–718.
- Miller, J. D. and Kent, D. V., 1988, Regional trends in the timing of Alleghenian remagnetization in the Appalachians: *Geology*, v. 16, p. 588–591.
- Needham, C. E. and Bates, R. L., 1943, Permian type sections in central New Mexico: Geological Society of America *Bulletin*, v. 54, p. 1653–1667.
- Onstott, T. C., 1980, Application of the Bingham distribution function in paleomagnetic studies: *Journal of Geophysical Research*, v. 85, p. 1500–1510.
- Pulliaiah, G. E., Irving, E., Buchan, K. L. and Dunlop, D. J., 1975, Magnetization changes caused by burial and uplift: *Earth and Planetary Science Letters*, v. 28, p. 133–143.
- Romer, A. S., 1960, The vertebrate fauna of the New Mexico Permian: New Mexico Geological Society, *Guidebook* 11, p. 48–54.
- Roy, J. L. and Park, J. K., 1974, The magnetization process of certain red beds: vector analysis of chemical and thermal results: *Canadian Journal of Earth Sciences*, v. 11, p. 437–471.
- Smith, R. L., Bailey, R. A. and Ross, C. S., 1970, Geologic map of the Jemez Mountains, New Mexico: U.S. Geological Survey, *Miscellaneous Geologic Investigations Map* I-571, scale 1:125,000.
- Starquist, V. L., 1988, Core log, Valles Caldera #2A, New Mexico: Los Alamos National Laboratory, Report LA-11176-OBES, 87 p.
- Steiner, M. B., 1988, Paleomagnetism of the Late Pennsylvanian and Permian: a test of the rotation of the Colorado Plateau: *Journal of Geophysical Research*, v. 93, p. 2201–2215.
- Van der Voo, R., 1989, Paleomagnetism of continental North America: the craton, its margins, and the Appalachian belt: Geological Society of America, *Memoir* 172, p. 447–470.
- Walton, D., 1980, Time-temperature relations in the magnetization of assemblies of single domain grains: *Nature*, v. 286, p. 245–247.
- WoldeGabriel, G. and Goff, F., 1989, Temporal relations of volcanism and hydrothermal systems in two areas of the Jemez volcanic field, New Mexico: *Geology*, v. 17, p. 986–989.
- WoldeGabriel, G. and Goff, F., 1992, K/Ar dates of hydrothermal clays from core hole VC-2B, Valles caldera, New Mexico and their relation to alteration in a large hydrothermal system: *Journal of Volcanology and Geothermal Research*, v. 50, p. 207–230.
- Zijderveld, J. D. A., 1967, A.C. demagnetization of rocks: analysis of results; *in* Collinson, D. W., Creer, K. M. and Runcorn, S. K., *Methods in palaeomagnetism*: Amsterdam, Elsevier, p. 254–286.