

New Mexico Geological Society

Downloaded from: <http://nmgs.nmt.edu/publications/guidebooks/6>



Some recent discoveries of uranium in Sierra County, New Mexico

Boyd, Fred S., Jr., 1955, pp. 123

in:
South-Central New Mexico, Fitzsimmons, J. P.; [ed.], New Mexico Geological Society 6th Annual Fall Field Conference Guidebook, 193 p.

This is one of many related papers that were included in the 1955 NMGS Fall Field Conference Guidebook.

Annual NMGS Fall Field Conference Guidebooks

Every fall since 1950, the New Mexico Geological Society (NMGS) has held an annual [Fall Field Conference](#) that explores some region of New Mexico (or surrounding states). Always well attended, these conferences provide a guidebook to participants. Besides detailed road logs, the guidebooks contain many well written, edited, and peer-reviewed geoscience papers. These books have set the national standard for geologic guidebooks and are an essential geologic reference for anyone working in or around New Mexico.

Free Downloads

NMGS has decided to make peer-reviewed papers from our Fall Field Conference guidebooks available for free download. Non-members will have access to guidebook papers two years after publication. Members have access to all papers. This is in keeping with our mission of promoting interest, research, and cooperation regarding geology in New Mexico. However, guidebook sales represent a significant proportion of our operating budget. Therefore, only *research papers* are available for download. *Road logs, mini-papers, maps, stratigraphic charts*, and other selected content are available only in the printed guidebooks.

Copyright Information

Publications of the New Mexico Geological Society, printed and electronic, are protected by the copyright laws of the United States. No material from the NMGS website, or printed and electronic publications, may be reprinted or redistributed without NMGS permission. Contact us for permission to reprint portions of any of our publications.

One printed copy of any materials from the NMGS website or our print and electronic publications may be made for individual use without our permission. Teachers and students may make unlimited copies for educational use. Any other use of these materials requires explicit permission.

This page is intentionally left blank to maintain order of facing pages.

SOME RECENT DISCOVERIES OF URANIUM IN SIERRA COUNTY, NEW MEXICO

By Fred S. Boyd, Jr.
*U. S. Atomic Energy Commission
Division of Raw Materials*

Increased prospecting activity in Sierra County, New Mexico, as in the rest of western United States, has resulted in the discovery of many new uranium occurrences. To date, no sizable tonnages of uranium ore have been developed in this area, but it is probable that at least a small production will be forthcoming from the more promising discoveries now known.

In eastern Sierra County on the east flank of the Caballo Mountains secondary uranium minerals are associated with secondary copper minerals and carbon trash in siltstones of the Abo formation of Permian age. Uranium minerals are most commonly found along vertical fractures and around carbon trash in small lenticular beds of light green siltstone. The size of these deposits varies from a few inches to eight feet in thickness and from a few feet to 40 or more feet in length. Uranium minerals commonly are spottily distributed within these zones.

Uranium minerals are found in a fissure vein of fluorite located 3 miles southeast of Truth or Consequences on the west side of the Caballo Mountains. The deposit is in a northwest-trending fault zone which separates Precambrian granite from the Bliss sandstone of Cambrian age. Purple fluorite with finely disseminated uraninite (?) and with minor uranophane and secondary copper minerals is in the vein adjacent to the Precambrian granite hanging wall.

A minor occurrence of brannerite, an oxide of uranium and titanium, has been found in a small shear zone in Precambrian schist and granite on the east side of the Rio Grande opposite Carrie Tingley Hospital in Truth or Consequences.

In southeastern Sierra County, uranium occurs locally in the Garfield fault zone four miles east of Derry. At this point, the fault separates Paleozoic sediments of the Montoya group from Paleozoic sediments of the Abo formation and the Magdalena group. Secondary uranium minerals, associated with limonite and calcite, coat fractures and cavities in a boxwork of quartz. Uranium minerals are sparsely distributed through the 10-to 20-foot wide fault zone at two widely separated points along the fault.

In western Sierra County, occurrences of secondary uranium minerals in the Abo formation, generally similar to those of eastern Sierra County, are found in the Sierra Cuchillo near the town of Chise and in the Iron Mountain District about seven miles northeast of Winston. The occurrences in the Sierra Cuchillo generally are present in small limestone lenses 1 to 2 feet in thickness and 10 to 15 feet in length. Uranium minerals present are usually secondary, with one notable exception, in which uraninite, closely associated with pyrite and chalcopyrite, is disseminated through a small lenticular zone in limestone.

Secondary uranium minerals, associated with secondary copper minerals, are found in silicified siltstone and limestone in moderately east-dipping beds near the Iron Mountain fault block. According to Jahns (1944) these beds represent the west flank of a syncline. A number of felsic Tertiary intrusives are closely associated with the occurrences, suggesting a possible genetic relationship.

Although no detailed study has yet been made, preliminary field and laboratory evidence suggests a probable hydrothermal origin and a Tertiary age for most of these deposits in Sierra County.

Based on the deposits thus far discovered, certain geologic conditions or environments appear to merit particular attention in future search for new deposits in Sierra County. Fault, fracture, or shear zones and the rocks adjoining them, particularly in or near established mining districts and/or near intermediate to felsic intrusives, should receive primary consideration in this area as favorable hosts for uranium deposits. New discoveries in this type of geologic environment are increasing in and along the Rio Grande trench in central and south-central New Mexico. Where the above-mentioned geologic conditions exist, further search of the Abo formation appears to be particularly warranted.

REFERENCES

- Jahns, R. H. (1944) Beryllium and tungsten deposits of the Iron Mountain district, Sierra and Socorro Counties, New Mexico: U. S. Geol. Survey Bull. 945-C, p. 51.
Kelley, V. C., and Silver, C. (1952) Geology of the Caballo Mountains: Univ. of New Mexico Pub. in Geol., No. 4.