

New Mexico Geological Society

Downloaded from: <http://nmgs.nmt.edu/publications/guidebooks/62>



A paleontological survey of a part of the Tesuque Formation near Chimaya, New Mexico, and a summary of the biostratigraphy of the Pojoaque Member (Middle Miocene, Late Barstovian)

Scott B. Aby, Gary S. Morgan, and Daniel J. Koning, 2011, pp. 347-358

in:

Geology of the Tusas Mountains and Ojo Caliente, Author Koning, Daniel J.; Karlstrom, Karl E.; Kelley, Shari A.; Lueth, Virgil W.; Aby, Scott B., New Mexico Geological Society 62nd Annual Fall Field Conference Guidebook, 418 p.

This is one of many related papers that were included in the 2011 NMGS Fall Field Conference Guidebook.

Annual NMGS Fall Field Conference Guidebooks

Every fall since 1950, the New Mexico Geological Society (NMGS) has held an annual [Fall Field Conference](#) that explores some region of New Mexico (or surrounding states). Always well attended, these conferences provide a guidebook to participants. Besides detailed road logs, the guidebooks contain many well written, edited, and peer-reviewed geoscience papers. These books have set the national standard for geologic guidebooks and are an essential geologic reference for anyone working in or around New Mexico.

Free Downloads

NMGS has decided to make peer-reviewed papers from our Fall Field Conference guidebooks available for free download. Non-members will have access to guidebook papers two years after publication. Members have access to all papers. This is in keeping with our mission of promoting interest, research, and cooperation regarding geology in New Mexico. However, guidebook sales represent a significant proportion of our operating budget. Therefore, only *research papers* are available for download. *Road logs, mini-papers, maps, stratigraphic charts*, and other selected content are available only in the printed guidebooks.

Copyright Information

Publications of the New Mexico Geological Society, printed and electronic, are protected by the copyright laws of the United States. No material from the NMGS website, or printed and electronic publications, may be reprinted or redistributed without NMGS permission. Contact us for permission to reprint portions of any of our publications.

One printed copy of any materials from the NMGS website or our print and electronic publications may be made for individual use without our permission. Teachers and students may make unlimited copies for educational use. Any other use of these materials requires explicit permission.

This page is intentionally left blank to maintain order of facing pages.

A PALEONTOLOGICAL SURVEY OF A PART OF THE TESUQUE FORMATION NEAR CHIMAYÓ, NEW MEXICO, AND A SUMMARY OF THE BIOSTRATIGRAPHY OF THE POJOAQUE MEMBER (MIDDLE MIocene, LATE BARSTOVIAN)

SCOTT B. ABY¹, GARY S. MORGAN², AND DANIEL J. KONING³

¹Muddy Spring Geology P.O. Box 488 Dixon, NM 87527; scottandlluvia@gmail.com

²New Mexico Museum of Natural History and Science, 1801 Mountain Rd., NW, Albuquerque, NM 87104

³New Mexico Bureau of Geology and Mineral Resources 801 Leroy Place Socorro, NM 87801

ABSTRACT—A recent paleontological survey north of Chimayó, NM, identified many new localities containing Miocene mammal fossils and one site containing freshwater(?) gastropods. All fossils were found within eight sections (T21N, R9E) at the southwest end of the survey area within the Pojoaque Member of the Tesuque Formation. The majority of the fossils were found within ~120 m stratigraphic interval containing the Pojoaque white ash zone, which to the south contains an ash that is 13.7 ± 0.18 Ma. Correlation of the Pojoaque white ash zone with the magnetic-polarity stratigraphy of Barghoorn, (1981), indicates an age range of 14.0-13.2 Ma for this ash zone. All fossils were found within sediments derived from Proterozoic metamorphic and Paleozoic sedimentary (sandstone, limestone, and shale) terrains to the northeast with minor and variable input of volcaniclastic material from the Latir volcanic field near Taos. Laterally adjacent and overlying sediments derived primarily from Proterozoic granite to the east are barren of fossil material. The fossil-bearing strata have been interpreted to represent a basin-floor environment while the barren, granitic sediments are interpreted as a piedmont environment. Preferential fossilization may reflect either different diagenetic conditions in the lime-bearing sediments relative to the sediments derived primarily from Proterozoic sources or reflect some inherent difference in depositional environment. A compilation of published paleontological data along with our recent findings produces a summary of the known fauna of the Pojoaque Member of the Tesuque Formation. Approximately 65 species of mammals are currently known from the Pojoaque Member, which represents one of the most important samples of early late Barstovian (early middle Miocene) mammals in western North America.

HISTORY OF STUDY AND OUTLINE OF STRATIGRAPHY

The Santa Fe Group (Spiegel and Baldwin, 1963) was extensively sampled for vertebrate remains by the American Museum of Natural History between 1924 and 1960 (Galusha and Blick, 1971). Less-extensive sampling was conducted as early as the 1870's by E.D. Cope and by the New Mexico Museum of Natural History since the 1970s. The American Museum collections include at least 20,000 "major skeletal elements", but only about 30% had been "formally catalogued" into their collection as of 1979 (Kues and Lucas, 1979, p. 238). No comprehensive assessment or description of these specimens has been conducted and little information on the location or stratigraphic context of the specimens is available in published form. What information is available is restricted to monographs or reports on individual taxa (see Kues and Lucas, 1979 and McKinney et al., 2001 for references).

The Santa Fe Group includes Oligocene-Quaternary basin-fill of the Española Basin of the Rio Grande rift (Spiegel and Baldwin, 1963). In the study area these rocks have been subdivided in two different ways. Galusha and Blick, (1971) defined a stratigraphy consisting of the Tesuque Formation with five members (Nambé, Skull Ridge, Pojoaque, Ojo Caliente Sandstone, and Chama-El Rito). These members were defined by stratigraphic markers that were based partially on biostratigraphy. The stratigraphic markers used by Galusha and Blick (1971) to define these members were not consistently found in the study area by Koning (2003) and Koning and Manley (2003) and they instead mapped 24 informal lithologic units. Koning (2003) did, however, map an approximate Pojoaque/Skull Ridge contact in the study area

(Fig. 1). The divisions of Koning (2003) and Koning and Manley (2003) were based on the work of Cavazza (1986), who defined two sedimentary provinces (A and B) with further subdivisions based on textural and bedding characteristics. Cavazza's provinces were defined by "sandstone and conglomerate petrology" (grain and clast type), paleocurrent data, and "sedimentary facies analysis". The sedimentary provinces of Cavazza (1986) are represented as two "lithosomes" (A and B) by Koning (2003) and Koning and Manley (2003).

In outcrop, these two lithosomes are most easily distinguished by grain and clast type. Gravels of lithosome A are dominated by granite, with lesser quartzite, and the sand fraction is arkosic (Cavazza, 1986; Koning, 2003; Koning and Manley, 2003). In the study area, gravels of lithosome B are mainly Paleozoic sandstone, siltstone, and limestone, with 5-20% Proterozoic quartzite, 0-20% felsic-intermediate volcanic rocks, and 1-5% granite (Koning et al., 2005). The sand of lithosome B is a litharenite to lithic arkose (Koning et al., 2005). Lithosome A has been interpreted as an alluvial slope or piedmont deposit laid down by west-flowing, intermittent streams (Smith, 2004; Kuhle and Smith, 2001), and lithosome B as basin-floor deposits laid down by a south-southwest-flowing river (Koning, et al., 2005).

PURPOSE AND METHODS OF THIS STUDY

In 2009-11 a paleontological survey was conducted on parts of the El Palacio/Fun Valley BLM off-road vehicle area located north of Chimayó and east of Ohkay Owingeh (formerly San Juan Pueblo), New Mexico. The objective of this study was to assess past impacts to fossil resources and limit any future impacts by identifying fossil-rich areas. While Gary Morgan and crews from




FIGURE 1. Geologic map of the fossil-rich part of the study area. Additional areas to the north and east of figure were surveyed but barren of fossil material. For additional information see Koning (2003) and Koning and Manley (2003). CS=Cuarceles stratigraphic section.

the NMMNH focused on collecting in known fossil-rich areas (Galusha and Blick, 1971; Kues and Lucas, 1979), Scott Aby surveyed all of BLM-managed parts of sections 5, 6, 7, 8, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 28, 30, 31, 32, 33 and 34 and

parts of sections 27 and 29, T21N, R9E (Fig. 1). The latter survey was conducted by ‘prospecting’ major and tributary drainages for fossil material, making transects perpendicular to strike of beds (usually along ridges or in drainages), and then thoroughly

searching any fossil-bearing horizons found in these transects. The location of all fossil sites was recorded using a GPS unit and their stratigraphic context was confirmed by field observations and comparison to geologic maps of the area (Koning, 2003; Koning and Manley, 2003). Fossils and fossil data are archived at the New Mexico Museum of Natural History in Albuquerque, NM.

FOSSIL DATA

Sedimentology

Fifty-six individual fossil localities were identified within the twenty-two sections (or parts of sections) surveyed by Aby. Some of these sites consist of multiple, closely spaced fossils. All of these sites were within the eight sections in the southwest part of the study area (within sections 27, 28, 29, 30, 31, 32, 33 and 34 of T21N, R9E; Fig. 1), an area already known to be fossil rich based on the work of the American Museum of Natural History (Appendix 1). Unidentifiable bone ‘shards’ or ‘chips’ were also found in section 20 of T21N, R9E.

Fifty-three of the fossil sites are in the Pojoaque Member of the Tesuque Formation and three are in the Cejita Member, but all are from ‘lithosome B’ sediments (e.g. sediments containing a mixture of Paleozoic sedimentary clasts and Proterozoic (plutonic and metamorphic) clasts ± Cenozoic volcanic clasts). Only five fossil sites are from areas that contained Cenozoic volcanic clasts as part of their pebble-sized component, and such beds are proportionately rare in the study area. It should be noted that the lithosome A or B designation was often made based on the nearest coarse-grained beds in the section, as fossils are commonly found in fine-grained beds. Spot checking confirmed that the pebble-sized fraction matched the general composition of the sand-sized fraction. Thirteen (~23%) of these sites contained material that was unambiguously “in place” (in their original location within the strata) and four more were probably in place. The remainder of the fossil material was found on the surface and may have been transported downslope, although in almost all cases this transport was probably limited to less than 100 m. Fourteen sites were located in or on green or greenish beds that we interpret as indicating reducing conditions and twenty-one sites were on or in red or reddish beds interpreted as indicating strongly oxidizing conditions. The remainder of sites were found in or on tan or brownish sediments interpreted as moderately-to-weakly oxidizing. Thirty-three out of fifty-six (59%) fossil sites were found in units interpreted by Koning (2003) as of ‘mixed’ provenance (i.e. interbedded lithosome A and B; Figs. 1, 2, Table 1). These sites were found in or on individual beds containing Paleozoic +/- volcanic clasts.

Age

All but one of Aby’s fossil sites are above Koning’s (2003) projected Pojoaque/Skull Ridge contact (Fig. 1) and are therefore from the Pojoaque Member of Galusha and Blick (1971), which contains fossils from the late Barstovian North American land

mammal “age” (NALMA; ~14.5-12 Ma; Tedford and Barghoorn, 1993; Tedford et al., 2004), consistent with the fossils found during our study (Appendix 1). Approximately eighty percent of the fossils we found are within ~1 km of the Pojoaque white ash zone (PWAZ), as mapped by Koning (2003) and Koning and Manley (2003; Fig. 2). The PWAZ is well-exposed 11 km south of here, in the Pojoaque Member type section of Galusha and Blick (1971). Here, one of the lower ashes of the PWAZ has been dated at 13.7 ± 0.18 Ma by the $^{40}\text{Ar}/^{39}\text{Ar}$ technique (Izett and Obradovich, 2001). Projection of the dated ash to the paleomagnetic section of Barghoorn (1981) (which was measured along the Pojoaque Member type section) allows us to constrain the age range of the Pojoaque white ash zone to 14.0-13.2 Ma. Chemical analyses of ashes in the study area indicates a match for two of these ashes with those in the Pojoaque Member type section (Slate et al., in review). The PWAZ in the Cuarteles measured section is approximately 120 m thick (Fig. 2). Using an average stratal accumulation rate of 0.15 mm/yr for the PWAZ in the study area (Koning et al., 2005), it would take approximately 0.8 Ma to deposit this much sediment.

Environmental Interpretation

At the time the fossils in the study area were deposited, the area was situated near the confluence of a broad, undissected alluvial-slope (lithosome A) and a fluvial basin floor (lithosome B) with highlands to the east and northeast and the Jemez Mountains volcanic field beginning to develop to the west (Kuhle and Smith, 2001, Smith, 2004). With time, the border between the basin floor and alluvial slope environments migrated back and forth across the study area.

Fossil flora that could be used to make local climate inferences for the Miocene in this area were long limited to a palm stump found in the Skull Ridge Member “...a few miles northwest of Santa Fe.” (Axelrod and Bailey, 1976). This palm fossil was used

TABLE 1. Representative fossils shown on Cuarteles Strat Section (Figure 2)

NMMNH number	Species	Description
28972	<i>Gomphotherium productum</i>	Mastodon
57608	<i>Meryceros crucensis</i>	Pronghorn
57620	<i>Aelurodon ferox</i>	Borophagine dog
63412	<i>Aelurodon ferox</i>	Borophagine dog
63413	<i>Pseudaelurus stouti</i>	Small cat
63415	<i>Pliogale nambiana</i>	Skunk
63416	<i>Copemys loxodon</i>	Rodent
63417	<i>Neohipparrison coloradense</i>	Horse
63418	<i>Merychys medius</i>	Oreodont
63419	<i>Merychys medius</i>	Oreodont
63420	<i>Hesperotestudo</i>	Giant land tortoise
L-7782	Various small vertebrates	Microfauna Quarry

Notes: Number preceded by an “L” is a locality number, all others are catalog numbers.




FIGURE 2. Cuardeles stratigraphic section with representative fossil sites projected onto it. Figure also shows correlation with Pojoaque Member type section and its dated ash. Age diagnostic fossils from these sites are listed in Table 1. Note that sites 63415 and 63412, although projecting into the lithosome A "Salmon Interval" of Galusha and Blick (1971,) were found in laterally adjacent lithosome B sediments.

to infer frost-free conditions. Recently, plant-fossil rich horizons have been found in the Pojoaque Member (McKinney, et al., 2006) that contain bristlecone pine and willow fossils among others (McKinney, personal comm., 2011). These new discoveries indicate that the climate varied from frost-free (Palm) to possibly “subalpine” (bristlecone pine) during deposition of the Tesuque Formation as a whole, or possibly that the drainage basin(s) contained this range of environments.

FOSSILIZATION

Clearly, bone was preferentially fossilized within sediments containing a mixture of Paleozoic and Proterozoic clasts (lithosome or province B) as opposed to those derived only from Proterozoic sources (lithosome or province A). The purpose of this study was simply to delineate areas that contain fossils, but we can offer some speculation on why bones were preferentially preserved in one type of sediment. Paleozoic clasts in the study area are dominated by sandstone with subordinate limestone, siltstone and mudstone clasts. Siltstone and mudstone are commonly ‘limey’ and sandstone commonly has some carbonate cement. We assume, therefore, that the groundwater moving through these sediments immediately after deposition had a higher pH than groundwater moving through the sediments of ‘lithosome A’. Lyman (1994) states that preservation of bone is better in high pH (alkaline) environments due, at least in part, to the inhibition of bacterial activity. Many fossils were found in sediments we interpret to represent oxidizing conditions (red, tan, and brownish sediments), so oxidizing conditions did not preclude preservation of bone in the study area. However, as green or greenish beds make up < 5% of the section, but accounted for ~25% of fossil sites, reducing conditions probably did favor bone preservation even within the context of overall good preservation potential.

The two types of sediment (A and B) have been interpreted as representing different depositional environments. Cavazza (1986) interpreted province A as an “...ephemeral environmental regime...” and province B as “...a more perennial regime.” (Cavazza, 1986, p. 287). This may have led to quicker burial of bone in lithosome B environments and therefore enhanced preservation potential. The abundance of fossils in sediments interpreted to represent interbedding of the two sediment types (Koning, 2003; Koning and Manley, 2003) may also reflect faster burial in these environments or some other factor unique to this depositional environment.

Many parts of the Tesuque Formation that contain abundant Paleozoic clasts do not contain fossils or are very sparsely fossiliferous; for example the Dixon and Cejita members to the north of the study area are barren (Koning and Aby, 2003; Manley, 1977, Morgan et al., this volume). By the same token, some highly fossiliferous parts of the Tesuque Formation contain no Paleozoic clasts (e.g., the Chama-El Rito Member near Medanales, NM; Koning et al., 2004). Factors unique to each fossiliferous part of the Santa Fe Group must therefore account for preservation in each area.

BIOSTRATIGRAPHY OF THE POJOAQUE MEMBER

The area surveyed during this study encompasses much of the Frick Laboratory “Santa Cruz fossil-collecting locality” (Galusha and Blick 1971, fig. 2; Fig. 1). Historically, this has been one of the richest areas for Miocene vertebrate fossils in the entire Española basin (e.g., Galusha and Blick, 1971; Tedford, 1981). The most productive of the Santa Cruz fossil sites are in the Pojoaque Member of the Tesuque Formation in Arroyo del Llano (First Wash of the Frick Laboratory) and Arroyo de Quarteles (Second Wash of the Frick Laboratory). East of Arroyo de la Morada (Third Wash of the Frick Laboratory), exposures are in either the Skull Ridge or Nambé Members of the Tesuque Formation. Only one fossil site was found in the Skull Ridge Member during our survey and none in the Nambé Member. Both the Skull Ridge and Nambé Members in this area (as mapped by Galusha and Blick, 1971) are composed mostly of “lithosome A” sediments (Koning, 2003) Only the fossils from the Pojoaque Member in the Santa Cruz area are discussed below.

Appendix 1 provides a faunal list of the late Barstovian (medial Miocene) mammals identified from the Pojoaque Member of the Tesuque Formation in the central Española basin, encompassing the Santa Cruz, North Pojoaque Bluffs, and Central Pojoaque Bluffs “fossil-collecting localities” of the Frick Laboratory (Galusha and Blick, 1971, fig. 2, localities 5, 6, 7). The late Barstovian mammal fauna from the Pojoaque Member reviewed by Tedford (1981) and Tedford and Barghoorn (1993) was referred to as the “Santa Cruz sites.” However, it is clear from the discussions in these papers that the fauna from the “Santa Cruz sites” was derived not only from the Santa Cruz fossil-collecting locality of Galusha and Blick (1971), as defined above, but also from the North Pojoaque Bluffs and Central Pojoaque Bluffs fossil-collecting localities of the Frick Laboratory. The Pojoaque Bluffs sites (Galusha and Blick, 1971) and the Jacona Microfossil Quarry (Chaney, 2009), as well as the original “Santa Fe marls” sites of Cope (1877), are in the general area of the type section of the Pojoaque Member in the vicinity of the Pojoaque Bluffs (Los Barrancos on the Española 7.5 minute quadrangle), on the Pojoaque and San Ildefonso pueblos and Jacona land grant, about 10-12 km south of the actual Santa Cruz sites.

Taxa in Appendix 1 followed by a superscript ¹ are represented by identifiable fossils collected in Arroyo del Llano and Arroyo de Quarteles during our recent BLM survey and are housed in the New Mexico Museum of Natural History (NMMNH) in Albuquerque. Taxa in Appendix 1 followed by a superscript ² are represented by published fossils from the Pojoaque Member in the Santa Cruz area, mostly collected by Frick Laboratory paleontologists (Joe Rak, John Blick, and Ted Galusha) from First Wash and Second Wash. Taxa followed by a superscript ³ are known from the Pojoaque Bluffs area (North Pojoaque Bluffs and Central Pojoaque Bluffs fossil-collecting localities of the Frick Laboratory), including the Jacona Microfossil Quarry (Chaney, 2009). Taxa in Appendix 1 lacking a superscript number are derived from the Pojoaque Member in the Española basin and are known from the Santa Cruz and/or the Pojoaque Bluffs areas; however,

we have not been able to confirm the exact locality data for these taxa from the literature. The taxa lacking superscripts are included in Appendix 1 based primarily on biostratigraphic studies of Miocene mammals from the Española basin (Tedford, 1981; Tedford and Barghoorn, 1993; Tedford et al., 2004) or on mammals reviewed in the various papers in Janis et al. (1998a, 2008). The early late Barstovian locality listed in Janis et al. (1998b, p. 635 and 2008, p. 710) as “SB32D. Pojoaque Member (inc. ... Pojoaque Bluffs Area, Santa Cruz Sites, and Jacona Microfossil Quarry)(Ba2)...” actually includes essentially all late Barstovian mammals known from the Pojoaque Member in the Española basin. Thus, for the mammals listed from locality “SB32D” in various papers in Janis et al. (1998a, 2008), we are not able to distinguish taxa listed as coming from the Santa Cruz sites from those derived from Pojoaque Bluffs and other sites in the Pojoaque Member. In some of the taxonomic papers listed below (other than in the Janis et al., 1998a, 2008 volumes), the specific localities for specimens are listed, and we were able to determine which specimens were collected in the Santa Cruz area.

In addition to our recent field survey, the vertebrate faunal list for the Pojoaque Member in the Santa Cruz area comes from a variety of published sources, primarily taxonomic works on various mammalian taxa, including: Erinaceidae (Rich, 1981), Rodentia (Korth and Chaney, 1999; Korth, 2002, 2008), Mustelidae (Baskin, 1998), Ursidae (Hunt, 1998a), Amphicyonidae (Hunt, 1998b), Canidae (Wang et al., 1999; Tedford et al., 2009), Felidae (Rothwell, 2003), Equidae (MacFadden, 1984, 1998), Rhinocerotidae (Prothero, 1998), Oreodontidae (Schultz and Falkenbach, 1941; Lander, 1998), Camelidae (Honey et al., 1998), Dromomerycidae (=Palaeomerycidae of some authors; Frick, 1937; Janis and Manning, 1998a; Prothero and Liter, 2008), Moschidae (Frick, 1937; Webb, 1998; Prothero, 2008), Antilocapridae (Frick, 1937; Janis and Manning, 1998b), and Gomphotheriidae (Frick, 1933; Osborn, 1936; Tobein, 1973; Lambert and Shoshani, 1998).

Paleontologists from the Frick Laboratory (Joe Rak, John Blick, Ted Galusha, and others) made almost yearly field surveys of Miocene vertebrates in the Española basin over 40 years from the mid 1920s to the mid 1960s (Galusha and Blick, 1971; Galusha, 1974). From the late 1970s to the present, field crews from the New Mexico Museum of Natural History and University of New Mexico have conducted periodic surveys of this same region (Kues and Lucas, 1979). Sediments of the Pojoaque Member of the Tesuque Formation in the central Española basin, including the Santa Cruz and Pojoaque Bluffs sites of the Frick Laboratory (Galusha and Blick, 1971), have produced a diverse mammalian fauna numbering at least 65 species (Appendix 1). The intensive field activity in the Española basin over a period of more than 85 years would suggest that the large mammal fauna of the Pojoaque Member is almost completely documented, and that few additional species remain to be discovered. Possibilities for additions to the fauna include uncommon groups currently unrepresented, including chalicotheres (Chalicotheriidae), tapirs (Tapiridae), peccaries (Tayassuidae), and protoceratids (Protoceratidae). Future additions to the faunal list will almost certainly be dominated by small mammals. Only a single diverse microfauna

has been reported from the Pojoaque Member, the Jacona Microfossil Quarry (Chaney, 2009). Several other microvertebrate sites are known but inadequately described or sampled, including the Pojoaque Bluffs, the “red layer” in Arroyo de Quartales (Second Wash), and the “green layer” in Arroyo de la Morada (Third Wash). Ongoing field studies of the Pojoaque Member by NMMNH crews are focusing on the recovery of additional microvertebrate faunas. Overall, nearly a third of the mammalian taxa on the faunal list are identified only to the genus level. In most cases, this results from the lack of modern taxonomic studies of these groups, not an absence of adequate fossil material.

The Pojoaque Member also has a rather diverse lower vertebrate fauna composed of amphibians (frogs, toads, and salamanders), reptiles (land tortoises, freshwater turtles, snakes, and lizards), and a variety of birds. Lower vertebrates are not included in Appendix 1 because there are only a limited number of taxonomic studies of these groups in the Española Basin, and thus very few taxa have been identified below the family level. Moreover, unlike mammals, most Miocene lower vertebrates are not age diagnostic. Among lower vertebrates from the Pojoaque Member, the most commonly encountered in the field are medium to large tortoises of the genus *Hesperotestudo*, some with shells nearly a meter in length (Sena and Thomas, 1989). Chaney (2009) listed a wide range of amphibians, reptiles, and birds from the Jacona Microfossil Quarry, most identified only to the family level, including frogs/toads, salamanders, anguid lizards, colubrid and viperid snakes, and at least seven birds, including ducks (Anatidae), hawks (Accipitridae), quail (Tetraoninae), rails (Rallidae), owls (Strigidae), woodpeckers (Picidae), and perching birds (Passeriformes).

The following is a brief summary of the mammalian biostratigraphy of the Pojoaque Member in the central Española Basin, based on data from our field work and biostratigraphic studies by Tedford (1981), Tedford and Barghoorn (1993, 1997), and Tedford et al. (2004). The most age-diagnostic mammal in the Pojoaque Member is the gomphotheriid proboscidean *Gomphotherium productum*. The first appearance of proboscideans in North America as immigrants from Eurasia has been termed the “Proboscidean Datum” and occurred in the early medial Miocene (early late Barstovian) between about 14.5 and 14.8 Ma (Woodburne and Swisher, 1995; Tedford et al., 2004). Although there are rare records of proboscideans from the late early Miocene (late Hemingfordian NALMA) and earliest middle Miocene (early Barstovian NALMA) elsewhere in North America (e.g., Prothero et al., 2008), no proboscidean fossils have ever been found in the late Hemingfordian Nambé Member or early Barstovian Skull Ridge Member in the Española Basin. *Gomphotherium productum* is common in the Pojoaque Member, including a complete skull collected in 1988 (Heckert et al., 2000) and the associated mandibles of this same individual (NMMNH 28972) collected more than 20 years later during our current survey, from about 100 m above the base of the member in Arroyo del Llano (Fig. 2). The first appearance of several other mammals defines the late Barstovian (Ba2), including the following mammals known from the Pojoaque Member: the erinaceid hedgehog *Untermannix*, the amphicyonid bear dog *Pseudocyon*, the mephitid carnivore

Pliogale, and the ocootonid lagomorphs *Hesperolagomys* and *Russellogus* (Rich, 1981; Hunt, 1998b; Tedford et al., 2004; Chaney, 2009). We collected a lower jaw of *Pliogale nambiana* (NMMNH 63415) during our survey, near the top of the Pojoaque Member just below the Cejita Member (Fig. 2). Other mammals known from the Pojoaque Member that also have their earliest occurrence in the late Barstovian include: the borophagine canids *Aelurodon ferox* and *A. stirtoni*, the three-toed horse *Neohipparrison coloradense*, the oreodont *Merychodus medius*, the camelid *Procamelus*, the moschid ruminant *Longirostromeryx*, and the antilocaprid *Ramoceros*. During our survey, we collected a mandible (NMMNH 63412) and lower first molar (m1) of *Aelurodon ferox* (NMMNH 57620), both also from high in the Pojoaque Member. A partial skull, lower jaws, and partial postcranial skeleton of *Neohipparrison coloradense* (NMMNH 63417), was found about 80 m above the base of the Pojoaque Member. Other mammals from the Pojoaque Member that are typical of, but not limited to, the late Barstovian, include: the primitive bear *Hemicyon (Plithocyon) ursinus*, the small felid *Pseudaelurus stouti*, the fox-like canid *Leptocyon vafer*, the beaver *Monosaulax pansus*, the cricetid rodent *Copemys loxodon*, the large anchitheriine horse *Megahippus mckennai*, the dromomerycid artiodactyl *Cranioceras teres*, and the antilocaprid *Meryceros crucensis*. We found mandibles of *Pseudaelurus stouti* (NMMNH 63413), *Leptocyon vafer* (NMMNH 63414), and *Copemys loxodon* (NMMNH 63416), and a horn core of *Meryceros crucensis* (NMMNH 57608) during our recent survey. The stratigraphic position of several of these species in the Pojoaque Member is indicated in Fig. 2. The fossil mammals from the Pojoaque Member represent one of the most important samples of early late Barstovian mammals in western North America, dating to the interval between about 14.5 and 13.3 Ma (Tedford, 1981; Tedford and Barghoorn, 1993; Tedford et al., 2004), which supports the age range of this unit indicated by the Pojoaque white ash zone (14.0–13.2 Ma) as discussed above.

It is always preferable to determine biostratigraphic intervals based on the presence of taxa rather than their absence because there are many factors besides age that may result in taxa being absent from a particular fauna, including paleoecology, taphonomy, biogeography, and others. Nonetheless, in very well-sampled geologic units in which hundreds, if not thousands of fossils have been collected, such as the Pojoaque and Skull Ridge Members, both the presence and absence of taxa have biostratigraphic significance. Thus, the lack of Proboscidea in the Skull Ridge Member almost certainly reflects a true absence of this group, especially considering the abundance of *Gomphotherium* in the overlying Pojoaque Member. Conversely, it is also significant that several age-diagnostic genera of mammals have their last appearance in the Skull Ridge Member (early Barstovian, Ba1) and are absent from the Pojoaque Member, including: the large amphicyonid *Amphicyon*, the anchitheriine horse *Anchitherium*, the oreodont *Brachycrus*, the youngest stenomylidine camel *Rakomylus*, and the dromomerycid artiodactyls *Dromomeryx* and *Rakomeryx*.

CONCLUSIONS

The southeastern part of the study area is very fossil rich. These fossils are all found within sediments containing a mixture of Paleozoic and Proterozoic clast types ± Cenozoic volcanic clasts. Sediments composed of only Proterozoic clast types did not yield any fossil material. The fossils are late Barstovian in age based on the concept of North American land mammal “ages” (Tedford and Barghoorn, 1993; Tedford et al., 2004) and the majority of the fossils are within the Pojoaque White Ash Zone, which is 13.2–14.0 Ma (Koning, 2003). Diagenetic conditions after burial likely had a significant effect on preservation of bone in the study area, but the specific depositional environment and rate of deposition may have also been significant factors.

ACKNOWLEDGMENTS

This survey was conducted under the auspices of the Bureau of Land Management and overseen by Patricia Hester and Paul Williams. Cordell Arellano, Louise Browne, Lorali Aby, and Lluvia Aby all helped SBA with fossil collection and many sites would not have been found without them. GSM thanks Paul Sealey, Warren Slade, Richard Franz, Larry Rinehart, and Dwayne Ulibarri for their help in the field. This paper was reviewed by Spencer Lucas and Sean Connell.

REFERENCES

- Axelrod, D.I., and Bailey, H.P., 1976, Tertiary vegetation, climate, and altitude of the Rio Grande Depression, New Mexico–Colorado: *Paleobiology*, v. 2, p. 235–254.
- Baskin, J. A., 1998, Chapter 9. Mustelidae; in Janis, C. M., Scott, K. M., and Jacobs, I. L., eds., *Evolution of Tertiary mammals of North America, Volume 1: Terrestrial carnivores, ungulates, and ungulate-like mammals*: Cambridge, Cambridge University Press, p. 152–173.
- Barghoorn, S., 1981, Magnetic-polarity stratigraphy of the Miocene type Tesuque Formation, Santa Fe Group, in the Espanola Valley, New Mexico: *Geological Society of America Bulletin*, v. 105, p. 1538–1551.
- Barghoorn, S. F., 1985, Magnetic polarity stratigraphy of the Tesuque Formation, Santa Fe Group in the Espanola Valley, New Mexico, with a taxonomic review of the fossil camels [PhD dissertation]: Columbia University, New York, 430 p.
- Cavazza, W., 1986, Miocene sediment dispersal in the central Espanola Basin, Rio Grande rift, New Mexico, USA: *Sedimentary Geology*, v. 51, p. 119–135.
- Chaney, D. S., 2009, A brief overview of the geology, taphonomy, and paleoenvironment of the Jacona Microfauna (Miocene, late Barstovian), Pojoaque Member, Tesuque Formation, north-central New Mexico: *Museum of Northern Arizona, Bulletin* 65, p. 185–196.
- Cope, E. D., 1877, Report upon the extinct Vertebrata obtained in New Mexico by parties of the expedition of 1874, in, Wheeler, G. M., *Report upon United States Geological Surveys West of the One Hundredth Meridian: Washington*, v. 4, pt. 2, p. 1–370.
- Frick, C., 1926, The Hemicyoninae and an American Tertiary bear: *Bulletin of the American Museum of Natural History*, v. 56, p. 1–119.
- Frick, C., 1933, New remains of trilophodont-tetrabelodont mastodons: *Bulletin of the American Museum of Natural History*, v. 59, p. 505–652.
- Frick, C., 1937, Horned ruminants of North America: *Bulletin of the American Museum of Natural History*, v. 69, p. 1–669.
- Galusha, T., 1974, Dating rocks of the Santa Fe Group: programs and problems: N.M. Geological Society, 25th Field Conference Guidebook, p. 283–286.

- Galusha, T., and Blick, J.C., 1971, Stratigraphy of the Santa Fe Group, New Mexico: Bulletin of the American Museum of Natural History, v. 144, 127 p.
- Heckert, A. B., Lucas, S. G., and Morgan, G. S., 2000, Specimens of *Gomphotherium* in the New Mexico Museum of Natural History and Science and the species-level taxonomy of North American *Gomphotherium*: New Mexico Museum of Natural History and Science, Bulletin 16, p. 187-194.
- Honey, J. G., Harrison, J. A., Prothero, D. R., and Stevens, M. S., 1998, Chapter 30. Camelidae; in Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., Evolution of Tertiary mammals of North America, Volume 1: Terrestrial carnivores, ungulates, and ungulate-like mammals: Cambridge, Cambridge University Press, p. 439-462.
- Hunt, R. M., Jr., 1998a, Chapter 10. Ursidae; in Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., Evolution of Tertiary mammals of North America, Volume 1: Terrestrial carnivores, ungulates, and ungulate-like mammals: Cambridge, Cambridge University Press, p. 174-195.
- Hunt, R. M., Jr., 1998b, Chapter 11. Amphicyonidae; in Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., Evolution of Tertiary mammals of North America, Volume 1: Terrestrial carnivores, ungulates, and ungulate-like mammals: Cambridge, Cambridge University Press, p. 196-227.
- Izett, G.A., and Obradovich, J.D., 2001, 40Ar/39Ar ages of Miocene tuffs in basin-fill deposits (Santa Fe Group, New Mexico, and Troublesome Formation, Colorado) of the Rio Grande rift system: The Mountain Geologist, v. 38, no. 2, p. 77-86.
- Janis, C. M. and Manning, E. 1998a, Chapter 32. Dromomerycidae; in Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., Evolution of Tertiary mammals of North America, Volume 1: Terrestrial carnivores, ungulates, and ungulate-like mammals: Cambridge, Cambridge University Press, p. 477-490.
- Janis, C. M. and Manning, E. 1998b, Chapter 33. Antilocapridae; in Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., Evolution of Tertiary mammals of North America, Volume 1: Terrestrial carnivores, ungulates, and ungulate like mammals: Cambridge, Cambridge University Press, p. 491-507.
- Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., 1998a, Evolution of Tertiary mammals of North America, Volume 1: terrestrial carnivores, ungulates, and ungulate-like mammals: Cambridge, Cambridge University Press, 691 p.
- Janis, C. M., Scott, K. M., and Jacobs, L. L., 1998b, Appendix I: Tertiary mammal localities; in Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., Evolution of Tertiary mammals of North America, Volume 1: terrestrial carnivores, ungulates, and ungulate-like mammals: Cambridge, Cambridge University Press, p. 625-654.
- Janis, C. M., Gunnell, G. F., and Uhen, M. D., eds., 2008, Evolution of Tertiary mammals of North America, Volume 2: small mammals, xenarthrans, and marine mammals: Cambridge, Cambridge University Press, 795 p.
- Koning, D.J., 2003, Geologic map of the Chimayo 7.5-minute quadrangle, Rio Arriba and Santa Fe counties, New Mexico: New Mexico Bureau of Geology and Mineral Resources, Open-file Geologic Map OF-GM-71, scale 1:24,000.
- Koning, D.J., and Aby, S.B., 2003, Geologic map of the Velarde Quadrangle, Rio Arriba County, New Mexico: New Mexico Bureau of Geology and Mineral Resources, Open-file Geologic Map 79, scale 1:24,000.
- Koning, D.J., and Manley, K., 2003, Geologic map of the San Juan Pueblo 7.5-minute quadrangle, Rio Arriba and Santa Fe counties, New Mexico: New Mexico Bureau of Geology and Mineral Resources, Open-file Geologic Map OF-GM-70, scale 1:24,000.
- Koning, D.J., May, J., Aby, S.B., and Horning, R., 2004, Preliminary geologic map of the Medanales 7.5 Minute quadrangle, Rio Arriba County, New Mexico: New Mexico Bureau of Geology and Mineral Resources, Open-file Geologic Map OF-GM-89, scale 1:24,000.
- Koning, D.K., Connell, S.D., Morgan, G.S., Peters, L., and McIntosh, W.C., 2005, Stratigraphy and depositional trends in the Santa Fe Group near Española, north central New Mexico: Tectonic and climatic implications: N.M. Geological Society, 56th Field Conference Guidebook, p. 237-257.
- Korth, W. W., 2002, Topotypic cranial material of the beaver *Monosaulax pansus* Cope (Rodentia, Castoridae): Paludicola, v. 4, p. 1-5.
- Korth, W. W., 2008, Two new pocket mice (Mammalia, Rodentia, Heteromyidae) from the Miocene of Nebraska and New Mexico and the early evolution of the subfamily Perognathinae: Geodiversitas, v. 30, p. 593-609.
- Korth, W. W. and Chaney, D. S., 1999, A new subfamily of geomyoid rodents (Mammalia) and a possible origin of Geomyidae: Journal of Paleontology, v. 73, p. 1191-1200.
- Kues, B., and Lucas, S., 1979, Summary of the paleontology of the Santa Fe Group (Mio-Pliocene), North-Central New Mexico: N.M. Geological Society, 30th Field Conference Guidebook, p. 237-241.
- Kuhle, A.J., and Smith, G.A., 2001, Alluvial-slope deposition of the Skull Ridge Member of the Tesuque Formation, Española Basin, New Mexico: New Mexico Geology, v. 23 p. 30-37.
- Lambert, W. D., and Shoshani, J., 1998, Chapter 43. Proboscidea; in Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., Evolution of Tertiary mammals of North America, Volume 1: Terrestrial carnivores, ungulates, and ungulate-like mammals: Cambridge, Cambridge University Press, p. 606-621.
- Lander, B., 1998, Chapter 27. Oreodontoidea; in Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., Evolution of Tertiary mammals of North America, Volume 1: Terrestrial carnivores, ungulates, and ungulate-like mammals: Cambridge, Cambridge University Press, p. 402-425.
- Lyman, R.L., 1994, Vertebrate Taphonomy, Cambridge University Press 524 p.
- MacFadden, B. J., 1984, Systematics and phylogeny of *Hipparrison*, *Neohipparrison*, *Nannippus*, and *Cormohipparrison* (Mammalia, Equidae) from the Miocene and Pliocene of the New World: Bulletin of the American Museum of Natural History, v. 171, p. 1-195.
- MacFadden, B. J., 1998, Chapter 37. Equidae; in Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., Evolution of Tertiary mammals of North America, Volume 1: Terrestrial carnivores, ungulates, and ungulatelike mammals: Cambridge, Cambridge University Press, p. 537-559.
- Manley, K., 1977, Geologic map of the Cejita Member (new name) of the Tesuque Formation, Española Basin, New Mexico: U.S. Geological Survey Miscellaneous Field Studies Map MF-877, scale 1:24000.
- McKinney, K.C., Chaney, D.S., and Maldonado, F., 2006, Preliminary report on the newly discovered Miocene flora of the Pojoaque Bluffs, Española Basin, New Mexico. [Abstract] Española Basin Technical Advisory Committee (EBTAG) 5th annual Española Basin Workshop Preliminary Abstracts, p. 34.
- McKinney, K.C., Tedford, R.H., Morgan, G.S., and Williamson, T.E., 2001, Fossil mammals of the Middle Rio Grande—Annotated dataset and bibliography of New Mexico vertebrate paleontology: U.S. Department of the Interior Open-File Report 01-023, CD ROM.
- Morgan, G., Aby, S.B., and Vogel, M., 2011, A new discovery of a skeleton of the horse *Protohippus*? and a summary of Miocene (Barstovian) fossil localities near Dixon, Rio Arriba County, New Mexico: N.M. Geological Society, 62nd Field Conference Guidebook p. 339-346.
- Osborn, H. F., 1936, Proboscidea, Vol. 1: New York, American Museum Press, 802 p.
- Prothero, D. R., 1998, Chapter 42, Rhinocerotidae; in Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., Evolution of Tertiary mammals of North America, Volume 1: Terrestrial carnivores, ungulates, and ungulate like mammals: Cambridge, Cambridge University Press, p. 595-605.
- Prothero, D. R., 2008, Systematics of the musk deer (Artiodactyla: Moschidae: Blastsomerycinae) from the Miocene of North America: New Mexico Museum of Natural History and Science, Bulletin 44, p. 207-223.
- Prothero, D. R. and Liter, M. R., 2008, Systematics of the dromomerycines and aletomerycines (Artiodactyla: Palaeomerycidae) from the Miocene and Pliocene of North America: New Mexico Museum of Natural History and Science, Bulletin 44, p. 273-298.
- Prothero, D. R., Davis, E. B., and Hopkins, S. S. B., 2008, Magnetic stratigraphy of the Massacre Lake Beds (late Hemingfordian, early Miocene), northwest Nevada, and the age of the “proboscidean datum” in North America: New Mexico Museum of Natural History and Science, Bulletin 44, p. 239-245.
- Rich, T. H. V., 1981, Origin and history of the Erinaceinae and Brachycerinae (Mammalia, Insectivora) in North America: Bulletin of the American Museum of Natural History, v. 171, p. 1-116.
- Rothwell, T., 2003, Phylogenetic systematics of North American *Pseudaelurus* (Carnivora: Felidae): American Museum Novitates, no. 3403, 64 p.
- Schultz, C. B. and Falkenbach, C. H., 1941, Ticholetinae, a new subfamily of oreodonts: Bulletin of the American Museum of Natural History, v. 79, p. 1-105.
- Sena, A., and Thomas, T., 1989, A giant land tortoise (*Geochelone*) from the Santa Fe Group (Miocene), near Española, New Mexico: New Mexico Journal of Science, v. 29, p. 69-78.
- Slate, J.L., Sarna-Wojciechski, A.M., Koning, D.J., Wan, E., Wahl, D.B., Connell, S.D., and Perkins, M.E., *in review*, Upper Neogene tephrochronologic correlations in the northern Rio Grande Rift, New Mexico, *in* Hudson, M., and Grauch, V.J.S., eds., New Perspectives on the Rio Grande rift: From Tectonics to Groundwater: Geological Society of America, Special Paper.

- Smith, G.A., 2004, Middle to late Cenozoic development of the Rio Grande rift and adjacent regions in northern New Mexico, in G. Mack and K. Giles, eds., Geology of New Mexico: New Mexico Geological Society, Special Publication 11, p. 331-358.
- Speigel, Z. and Baldwin, B., 1963, Geology and water resources of the Santa Fe Area, New Mexico: United States Geological Survey, Water Supply Paper 1525, 258 p.
- Taylor, B. E. and Webb, S. D., 1976, Miocene Leptomerycidae (Artiodactyla, Ruminantia) and their relationships: American Museum Novitates, num. 2596, 1-22.
- Tedford, R. H., 1981, Mammalian biochronology of the late Cenozoic basins of New Mexico: Geological Society of America Bulletin, v. 92, p. 1008-1022.
- Tedford, R. H. and Barghoorn, S., 1993, Neogene stratigraphy and mammalian biochronology of the Española Basin, northern New Mexico: New Mexico Museum of Natural History and Science, Bulletin 2, p. 158-168.
- Tedford, R. H. and Barghoorn, S., 1997, Miocene mammals of the Española and Albuquerque basins, north-central New Mexico: New Mexico Museum of Natural History and Science, Bulletin 11, p. 77-95.
- Tedford, R. H., Wang, X., and Taylor, B. E., 2009, Phylogenetic systematics of the North American fossil Caninae (Carnivora: Canidae): Bulletin of the American Museum of Natural History, no. 325, 218 pp.
- Tedford, R. H., Albright, L. B., III, Barnosky, A. D., Ferrusquia-Villafranca, I., Hunt, R. M., Jr., Storer, J. E., Swisher, C. C., III, Voorhies, M. R., Webb, S. D., and Whistler, D. P., 2004, Mammalian biochronology of the Arikareean through Hemphillian interval (late Oligocene through early Pliocene epochs), in Woodburne, M. O., ed., Late Cretaceous and Cenozoic mammals of North America: biostratigraphy and geochronology: New York, Columbia University Press, p. 169-231.
- Tobien, H., 1973, On the evolution of mastodonts (Proboscidea, Mammalia), part 1: The bunodont trilophodont groups: Notizblatt des Hessischen Landesamtes für Bodenforschung zu Wiesbaden, v. 101, p. 202-276.
- Wang, X., Tedford, R. H., and Taylor, B. E., 1999, Phylogenetic systematics of the Borophaginae (Carnivora: Canidae): Bulletin of the American Museum of Natural History, no. 243, 391 p.
- Webb, S. D., 1998, Chapter 31, Hornless ruminants; in Janis, C. M., Scott, K. M., and Jacobs, L. L., eds., Evolution of Tertiary mammals of North America, Volume 1: Terrestrial carnivores, ungulates, and ungulate-like mammals: Cambridge, Cambridge University Press, p. 463-476.
- White, J. A., 1987, The Archaeolaginae (Mammalia, Lagomorpha) of North America, excluding *Archaeolagus* and *Panolax*: Journal of Vertebrate Paleontology, v. 7, p. 425-450.
- Woodburne, M. O. and Swisher, C. C., III, 1995, Land mammal high-resolution geochronology, intercontinental overland dispersals, sea level, climate, and vicariance; in Geochronology, time scales, and global stratigraphic correlation: Society of Economic Paleontologists and Mineralogists, Special Publication, no. 54, p. 335-364.

APPENDIX 1. Mammalian faunal list from the middle Miocene (late Barstovian) Pojoaque Member, Tesuque Formation, Española basin, northern New Mexico.

Mammalian Fauna

Lipotyphla

Erinaceidae

- Metechinus amplior*²
- Untermannerix copiosus*^{2,3}

Talpidae

- Achlyoscapter*?³
- Gaillardia* sp.³
- Mystipterus* sp.³
- Scapanoscapter*?^{1,3}

Soricidae

- Adeloblarina* sp.³
- Alluvisorex* sp.³
- Limnoecus* sp.³

Rodentia

Mylagaulidae

- Mylagaulus laevis*³

Sciuridae

- Tamias* sp. ^{1,3}
- Spermophilus* sp. ^{1,3}
- cf. *Petauristodon* sp.

Heteromyidae

- Mioperognathus willardi*³

Geomyoidea

Subfamily Mojavemyinae

- Mojavemys galushai*³
- Phelosacomys neomexicanus*³

Castoridae

- Monosaulax pansus*^{1,2,3}

Zapodidae

- Plesiosminthus* sp.³

Cricetidae

- Copemys loxodon*^{1,3}

References

- Rich (1981)
Rich (1981)

- Chaney (2009)
Chaney (2009)
Chaney (2009)
Chaney (2009)

- Chaney (2009)
Chaney (2009)
Chaney (2009)

- Chaney (2009)

- Chaney (2009)
Chaney (2009)
Chaney (2009)

- Korth (2008)

- Korth & Chaney (1999)
Korth & Chaney (1999)

- Cope (1877), Korth (2002), Chaney (2009)

- Chaney (2009)

- Cope (1877), Chaney (2009)

APPENDIX 1. (cont.)

Mammalian Fauna

Lagomorpha

Ochotonidae

Hesperolagomys sp.³
Russellagus sp.³

Leporidae

Hypolagus sp.^{1,3}
*Panolax sanctae fidei*³

Carnivora

Procyonidae

Bassariscus sp.³

Mustelidae

Pliogale nambiana^{1,3}

Ursidae

Plithocyon (= *Hemicyon*) *ursinus*³

Amphicyonidae

Pseudocyon sp.³

Canidae

Borophaginae

Aelurodon ferox^{1,2,3}
*Aelurodon stirtoni*²
Carpocyon webbi^{2,3}
*Epicyon saevus*²
Paratomarctus temerarius^{2,3}

Caninae

Leptocyon vafer^{1,2,3}

Felidae

Pseudaelurus marshi^{1,2}
Pseudaelurus stouti^{1,2}

Perissodactyla

Equidae

Anchitheriinae

Hypohippus sp.
Megahippus mckennai

Equinae: Hipparrisonini

Cormohipparion sp.
“*Merychippus*” *calamarius*
Neohipparrison *affine*
Neohipparrison *coloradense*¹

Equinae: Equini

Pliohippus supremus

Rhinocerotidae

Aphelops megalodus
Peraceras hessei
Peraceras projectum
Peraceras superciliosum
Teleoceras new sp.

Artiodactyla

Oreodontidae

Merychyus medius^{1,2,3}

Camelidae

Miolabinae

Miolabis sp.
Nothotylopus sp.
Paramiolabis sp.

Protolabinae

Michenia sp.
Protolabis sp.

References

- Chaney (2009)
Chaney (2009)
- Chaney (2009)
Cope (1877), White, (1987), Chaney (2009)
- Chaney (2009)
- Chaney (2009)
- Chaney (2009)
- Cope (1877), Baskin (1998)
- Frick (1926), Hunt (1998a)
- Hunt (1998b)
- Wang, Tedford & Taylor (1999)
Wang, Tedford & Taylor (1999)
- Tedford, Wang & Taylor (2009)
- Rothwell (2003)
Rothwell (2003)
- MacFadden (1998)
MacFadden (1998)
- MacFadden (1998)
MacFadden (1998)
MacFadden (1998)
MacFadden (1998)
- MacFadden (1998)
- Prothero (1998)
Prothero (1998)
Prothero (1998)
Prothero (1998)
Prothero (1998)
- Schultz & Falkenbach (1941), Lander (1998)
- Barghoorn (1985), Honey et al. (1998)
Barghoorn (1985), Honey et al. (1998)
Barghoorn (1985), Honey et al. (1998)
- Barghoorn (1985), Honey et al. (1998)
Barghoorn (1985), Honey et al. (1998)


APPENDIX 1. (cont.)

Camelinae: Camelini	
<i>Procamelus</i> sp.	Barghoorn (1985), Honey et al. (1998)
Camelinae: Lamini	
<i>Aepycamelus</i> sp.	Barghoorn (1985), Honey et al. (1998)
Camelinae: incertae sedis	
<i>Australocamelus</i> sp.	Barghoorn (1985), Honey et al. (1998)
Leptomerycidae	
<i>Pseudoparablastomeryx francescita</i>	Frick (1937), Taylor & Webb (1976), Webb (1998)
Moschidae	
<i>Blastomeryx francesca</i>	Frick (1937), Webb (1998), Prothero (2008)
<i>Longirostromeryx blicki</i> ^{2,3}	Frick (1937), Webb (1998), Prothero (2008)
Dromomerycidae (=Palaeomerycidae)	
<i>Cranioceras teres</i> ^{2,3}	Cope (1877), Frick (1937), Janis & Manning (1998a), Prothero and Liter (2008)
Antilocapridae	
<i>Meryceros crucensis</i> ^{1,2,3}	Frick (1937), Janis & Manning (1998b)
<i>Meryceros major</i>	Frick (1937), Janis & Manning (1998b)
<i>Plioceros blicki</i>	Frick (1937), Janis & Manning (1998b)
<i>Ramoceros marthae</i>	Frick (1937), Janis & Manning (1998b)
<i>Ramoceros ramosus</i> ^{2,3}	Frick (1937), Janis & Manning (1998b)
<i>Submeryceros crucianus</i> ²	Frick (1937), Janis & Manning (1998b)
Proboscidea	
Gomphotheriidae	
<i>Gomphotherium productum</i> ^{1,2,3}	Cope (1877), Tobien (1973)
<i>Megabelodon joraki</i>	Frick (1933), Lambert & Shoshani (1998)

Notes:

Only taxa identified to the genus or species level are included on this list. Taxa followed by a superscript (1) were found in Arroyo del Llano and/or Arroyo de Quarteles during our recent survey. Taxa followed by a superscript (2) were collected by the Frick Laboratory from the Santa Cruz fossil-collecting locality, mostly from First Wash (=Arroyo del Llano) and Second Wash (=Arroyo de Quarteles). Taxa followed by a superscript (3) are known from the Pojoaque Bluffs area (North Pojoaque Bluffs and Central Pojoaque Bluffs fossil-collecting localities of the Frick Laboratory), including the Jacona Microfossil Quarry (Chaney, 2009). Unmarked taxa are derived from the Pojoaque Member in the Española basin and are known from the Santa Cruz and/or the Pojoaque Bluffs areas; however, we have not been able to confirm the exact locality data for these taxa from the literature (e.g., Tedford, 1981; Tedford and Barghoorn, 1993; various papers in Janis et al., eds., 1998a). References for the identifications of mammals from the Pojoaque Member are listed for each genus/species.

AN ODE TO KYANITE



*It is possible that I am pushing through solid rock,
like a vein of ore, encased.*

*And I am so far in, I see no end,
and no distance: everything is close to my face,
and everything close to my face is stone.*

*I am not yet wise in grief-
so this great darkness makes me small.
But if it's you: make yourself fierce, break in:
Then your great transforming will happen to me,
and to you, the fullness of my cry.*

- Rainer Maria Rilke

mary simmons